

169

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



A3

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07H 21/04, C12Q 1/68, A61K 48/00		A1	(11) International Publication Number: WO 00/20435
			(43) International Publication Date: 13 April 2000 (13.04.00)
(21) International Application Number: PCT/US99/23171			(74) Agents: LICATA, Jane, Massey et al.; Law Offices of Jane Massey Licata, 66 E. Main Street, Marlton, NJ 00853 (US).
(22) International Filing Date: 5 October 1999 (05.10.99)			
(30) Priority Data: 09/167,109 6 October 1998 (06.10.98) US			
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/167,109 (CIP) Filed on 6 October 1998 (06.10.98)			
(71) Applicant (for all designated States except US): ISIS PHARMACEUTICALS, INC. [US/US]; 2292 Faraday Avenue, Carlsbad, CA 92008 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): BAKER, Brenda, F. [US/US]; 2147 Avenida Toronja, Carlsbad, CA 92009 (US). COWSERT, Lex, M. [US/US]; 3008 Newshire Street, Carlsbad, CA 92008 (US). MONIA, Brett, P. [US/US]; 7605 Nueva Castilla Way, La Costa, CA 92009 (US). XU, Xiaoxing, S. [CN/US]; 18 Main Street, #3, Madison, NJ 07940 (US).			(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(54) Title: ANTISENSE MODULATION OF EXPRESSION OF TUMOR NECROSIS FACTOR RECEPTOR-ASSOCIATED FACTORS (TRAFs)			
(57) Abstract Compositions and methods are provided for modulating the expression of tumor necrosis factor receptor-associated factor (TRAF). Antisense compounds, particularly antisense oligonucleotides, targeted to nucleic acids encoding TRAF are preferred. Methods of using these compounds for modulation of TRAF expression and for treatment of diseases associated with expression of TRAF are provided.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

-1-

ANTISENSE MODULATION OF EXPRESSION OF TUMOR NECROSIS FACTOR
RECEPTOR-ASSOCIATED FACTORS (TRAFs)

FIELD OF THE INVENTION

5 The present invention provides compositions and
methods for modulating the expression of tumor necrosis
factor receptor-associated factors (TRAFs). In particular,
this invention relates to antisense compounds, particularly
10 oligonucleotides, specifically hybridizable with nucleic
acids encoding human TRAFs. Such oligonucleotides have
been shown to modulate the expression of TRAFs.

BACKGROUND OF THE INVENTION

Tumor necrosis factor (TNF) receptor superfamily
members regulate cellular proliferation, differentiation
15 and apoptosis in inflammatory and immune responses. This
receptor superfamily comprises a group of related cell-
surface receptors including, but not limited to, types 1
and 2 TNF receptors (TNFR1 and TNFR2), Fas, CD27, 4-1BB,
CD40 and CD30. Signaling through TNF receptor superfamily
20 members is initiated by oligomerization of the receptors
with trimeric ligands, bringing intracellular domains in
close proximity (Pullen et al., *Biochemistry* 1998, 37,
11836-11845). Two families of adaptor proteins that
associate with TNF receptor superfamily members have been
25 identified: the TNF receptor-associated factor (TRAF)
family, and the death domain-containing protein family.

Members of the TRAF family of proteins share an amino-
terminal RING finger motif and a homologous carboxy-
terminal region, referred to in the art as the TRAF domain
30 (Yuan, J., *Curr. Opin. Cell Biol.* 1997, 9, 247-251. This
conserved carboxy-terminal region binds to receptor
cytoplasmic domains and mediates interactions with the
signaling proteins NF- κ B inducing kinase (NIK) and I-
TRAF/TANK (Cheng et al., *Science* 1995, 267, 1494-1498;

-2-

Cheng, G. and Baltimore, D., *Genes Dev.* 1996, 10, 963-973; Rothe et al., *Proc. Natl Acad. Sci. USA* 1996, 93, 8241-8246; Malinin et al., *Nature* 1997, 385, 540-544). A predicted coiled-coil region mediating TRAF homo- and hetero-oligomerization is in a less conserved region N-terminal to the TRAF domain (Cao et al., *Nature* 1996, 383, 443-446; Cheng et al., *Science* 1995, 267, 1494-1498; Rothe et al., *Cell* 1994, 78, 681-692; Sato et al., *FEBS Lett* 1995, 358, 113-118; and Takeuchi et al., *J. Biol. Chem* 1996, 271, 19935-19942).

The mammalian TRAF family currently includes six members, TRAF-1, TRAF-2, TRAF-3, TRAF-4, TRAF-5 and TRAF-6. These proteins have generally been found within the cytosols of cells, either in association with cytosolic vesicles or at the plasma membrane after addition of selected TNF family cytokines to the cells. Members of the TRAF family mediate signals for various different receptors. Subsets of TRAF family members have been shown to interact with the TNF receptor family members (TNFR2, CD40, CD30, LT β R, ATAR, OX40 and 4-1BB).

For example, TRAF-1 and TRAF-2 were identified by their ability to interact with the cytoplasmic domains of TNFR2 (Rothe et al., *Cell* 1994, 78, 681-691). TNFR2 has been associated with TNF's ability to stimulate cell proliferation and activation of NF κ B (Tartaglia et al., *Proc. Natl Acad. Sci. USA* 1991, 88, 9292-9296). TRAF-1 is believed to be involved in the regulation of apoptosis (Speiser et al., *J. Exp. Med.* 1997, 185, 1777-1783). Depletion of TRAF-2 and its co-associated proteins has also been shown to increase the sensitivity of the cell to undergo apoptosis during activation of death inducing receptors such as TNFR1 (Duckett, C.S. and Thompson, C.B., *Genes & Development* 1997, 11, 2810-2821; Yeh et al., *Immunity* 1997, 7, 715-725). Accordingly, the rate of

-3-

receptor-mediated TRAF-2 consumption and TRAF-2 translation has been suggested to play a dynamic role in the regulation of cell survival (Duckett, C.S. and Thompson, C.B., *Genes & Development* 1997, 11, 2810-2821). Targeted disruption of

5 the TRAF-2 gene in mice has also been shown to generate severe defects in c-Jun N-terminal kinase (JNK) activation through TNFR1 (Yeh et al., *Immunity* 1997, 7, 715-725).

TRAF-2 (Rothe et al., *Science* 1995, 269, 1424-1427), TRAF-3 (Cheng et al., *Science* 1995, 267, 1494-1498), TRAF-5

10 (Ishida et al., *Proc. Natl Acad. Sci USA* 1996, 93, 9437-9442) and TRAF-6 (Pullen et al., *Biochemistry* 1998, 37, 11836-11845) have also been shown to interact with the B lymphocyte receptor CD40. CD40 is a TNF receptor superfamily member that provides activation signals in

15 antigen presenting cells such as B cells, macrophages and dendritic cells. Activation of CD40 leads to B-cell survival, growth and differentiation. In 293T cells, expression of TRAF-3 suppressed constitutive activity of NFkB, whereas expression of TRAF-5 induced NFkB activity.

20 Targeted disruption of the TRAF-3 gene in mice causes impaired immune responses to T-dependent antigens and results in early postnatal lethality (Xu et al., *Immunity* 1996, 5, 407-415). TRAF-2, TRAF-5 or TRAF-6 overexpression in mammalian cells also induces JNK activation.

25 TRAF-4 is expressed in breast cancers. In in vitro binding assays, TRAF-4 has been shown to interact with the cytosolic domain of the lymphotoxin- β receptor (LT β R) and weakly with the p75 nerve growth factor receptor but not with TNFR1, TNFR2, Fas or CD40 (Karjewska et al., *Am. J. of*

30 *Pathol.* 1998, 152, 6, 1549-1561).

TRAF-6 has also been reported to mediate the signal transduction pathway induced by IL-1 to activate NFkB by recruiting IL-1 receptor associated kinase (IRAK), a serine/threonine kinase (Cao et al., *Nature* 1996 381:9437-

-4-

9442). Thus, the role of TRAFs extends beyond being signal transducers for the TNF-receptor superfamily.

The TRAF-5 protein and DNA encoding TRAF-5 are disclosed in WO97/38099. Also disclosed in WO97/38099 is
5 an antisense oligonucleotide against the DNA, an anti-TRAF-5 antibody, a vector containing the DNA, transformants containing this vector and methods of producing TRAF-5 with this vector. In addition, this PCT application discloses methods of screening substances binding to TRAF-5 and
10 substances regulating the activity and expression of this protein.

A TRAF family molecule, a polynucleotide coding for this molecule, an antibody against the molecule and an antisense polynucleotide of the molecule are also disclosed
15 in WO97/31110. Disclosed in this PCT application are the base sequence of the gene and the amino acid of this "unknown" TRAF family molecule, which in addition to the antibody, are suggested to provide means for elucidating the functions of the proteins and the signal transducer
20 system of a TNF-R family in which this molecule participates, to provide probes for research and diagnosis, and to indicate applications for therapeutic agents.

Currently, however, there are no known therapeutic agents which effectively inhibit the synthesis of one or
25 more selected TRAF family members. Consequently, there is a long-felt need for agents capable of effectively inhibiting TRAF expression. Antisense oligonucleotides against one or more TRAFs may therefore prove to be uniquely useful in a number of therapeutic, diagnostic and research
30 applications.

SUMMARY OF THE INVENTION

The present invention is directed to antisense compounds, particularly oligonucleotides, which are targeted to a nucleic acid encoding a selected tumor
35 necrosis factor receptor-associated factor (TRAF), and

-5-

which modulate the expression of the selected TRAF. Pharmaceutical and other compositions comprising the antisense compounds of the invention are also provided. Further provided are methods of modulating the expression of TRAFs in cells or tissues comprising contacting said cells or tissues with one or more of the antisense compounds or compositions of the invention. Further provided are methods of treating an animal, particularly a human, suspected of having or being prone to a disease or condition associated with expression of a selected TRAF by administering a therapeutically or prophylactically effective amount of one or more of the antisense compounds or compositions of the invention.

DETAILED DESCRIPTION OF THE INVENTION

The present invention employs oligomeric antisense compounds, particularly oligonucleotides, for use in modulating the function of nucleic acid molecules encoding selected tumor necrosis factor receptor-associated factors (TRAFs), ultimately modulating the amount of the selected TRAF produced. This is accomplished by providing antisense compounds which specifically hybridize with one or more nucleic acids encoding the selected TRAF. By "selected TRAF" it is meant any member of the TRAF family of proteins, most preferably TRAF-1, TRAF-2, TRAF-3, TRAF-4, TRAF-5 or TRAF-6. As used herein, the terms "target nucleic acid" and "nucleic acid encoding TRAF" encompass DNA encoding a TRAF family member, RNA (including pre-mRNA and mRNA) transcribed from such DNA, and also cDNA derived from such RNA. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal function of the nucleic acid. This modulation of function of a target nucleic acid by compounds which specifically hybridize to it is generally referred to as "antisense". The functions of DNA to be interfered with include replication and transcription. The functions of

-6-

RNA to be interfered with include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of the selected TRAF. In the context of the present invention, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of a gene. In the context of the present invention, inhibition is the preferred form of modulation of gene expression and mRNA is a preferred target.

It is preferred to target specific nucleic acids for antisense. "Targeting" an antisense compound to a particular nucleic acid, in the context of this invention, is a multi-step process. The process usually begins with the identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the present invention, the target is one or more nucleic acid molecules encoding one or more selected TRAFs. The targeting process also includes determination of a site or sites within this gene for the antisense interaction to occur such that the desired effect, e.g., detection or modulation of expression of the protein, will result. Within the context of the present invention, a preferred intragenic site is the region encompassing the translation initiation or termination codon of the open reading frame (ORF) of the gene. Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation

-7-

codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function *in vivo*. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used *in vivo* to initiate translation of an mRNA molecule transcribed from a gene encoding a TRAF, regardless of the sequence(s) of such codons.

It is also known in the art that a translation termination codon (or "stop codon") of a gene may have one of three sequences, *i.e.*, 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively). The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (*i.e.*, 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (*i.e.*, 5' or 3') from a translation termination codon.

The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the

-8-

translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene, and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap. The 5' cap region may also be a preferred target region.

Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as "introns", which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. mRNA splice sites, i.e., intron-exon junctions, may also be preferred target regions, and are particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular mRNA splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions are also preferred targets. It has also been found that introns can also be effective, and therefore preferred, target regions for antisense compounds targeted, for example, to DNA or pre-mRNA.

Once one or more target sites have been identified,

-9-

oligonucleotides are chosen which are sufficiently complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect.

- 5 In the context of this invention, "hybridization" means hydrogen bonding, which may be Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleoside or nucleotide bases. For example, adenine and thymine are complementary nucleobases which
- 10 pair through the formation of hydrogen bonds. "Complementary," as used herein, refers to the capacity for precise pairing between two nucleotides. For example, if a nucleotide at a certain position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the same
- 15 position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered to be complementary to each other at that position. The oligonucleotide and the DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each
- 20 molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically hybridizable" and "complementary" are terms which are used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the
- 25 oligonucleotide and the DNA or RNA target. It is understood in the art that the sequence of an antisense compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. An antisense compound is specifically hybridizable when
- 30 binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under
- 35 conditions in which specific binding is desired, i.e.,

-10-

under physiological conditions in the case of *in vivo* assays or therapeutic treatment, or in the case of *in vitro* assays, under conditions in which the assays are performed.

Antisense compounds are commonly used as research
5 reagents and diagnostics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes. Antisense compounds are also used, for example, to
10 distinguish between functions of various members of a biological pathway. Antisense modulation has, therefore, been harnessed for research use.

The specificity and sensitivity of antisense is also harnessed by those of skill in the art for therapeutic
15 uses. Antisense oligonucleotides have been employed as therapeutic moieties in the treatment of disease states in animals and man. Antisense oligonucleotides have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus
20 established that oligonucleotides can be useful therapeutic modalities that can be configured to be useful in treatment regimens in cells, tissues and animals, especially humans.

In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of
25 ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. This term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent internucleoside (backbone) linkages as well as oligonucleotides having non-naturally-occurring portions
30 which function similarly. Such modified or substituted oligonucleotides are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target and increased stability in the presence of
35 nucleases.

-11-

While antisense oligonucleotides are a preferred form of antisense compound, the present invention comprehends other oligomeric antisense compounds, including but not limited to oligonucleotide mimetics such as are described below. The antisense compounds in accordance with this invention preferably comprise from about 8 to about 30 nucleobases. Particularly preferred are antisense oligonucleotides comprising from about 8 to about 30 nucleobases (i.e., from about 8 to about 30 linked nucleosides). As is known in the art, a nucleoside is a base-sugar combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such heterocyclic bases are the purines and the pyrimidines. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to form a linear polymeric compound. In turn the respective ends of this linear polymeric structure can be further joined to form a circular structure, however, open linear structures are generally preferred. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA and DNA is a 3' to 5' phosphodiester linkage.

Specific examples of preferred antisense compounds useful in this invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the

-12-

backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

5 Preferred modified oligonucleotide backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkyl-phosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates and chiral phosphonates, 10 phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and 15 those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also included.

Representative United States patents that teach the 20 preparation of the above phosphorus-containing linkages include, but are not limited to, U.S.: 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 25 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050, each of which is herein incorporated by reference.

Preferred modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that 30 are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages 35 (formed in part from the sugar portion of a nucleoside);

-13-

siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH₂ component parts.

Representative United States patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S.: 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; and 5,677,439, each of which is herein incorporated by reference.

In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S.: 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching

-14-

of PNA compounds can be found in Nielsen et al., *Science*, 1991, 254, 1497-1500.

Most preferred embodiments of the invention are oligonucleotides with phosphorothioate backbones and
5 oligonucleosides with heteroatom backbones, and in particular -CH₂-NH-O-CH₂-, -CH₂-N(CH₃)-O-CH₂- [known as a methylene (methylimino) or MMI backbone], -CH₂-O-N(CH₃)-CH₂-, -CH₂-N(CH₃)-N(CH₃)-CH₂- and -O-N(CH₃)-CH₂-CH₂- [wherein the native phosphodiester backbone is represented as
10 -O-P-O-CH₂-] of the above referenced U.S. Patent 5,489,677, and the amide backbones of the above referenced U.S. Patent 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced U.S. Patent 5,034,506.

15 Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and
20 alkynyl may be substituted or unsubstituted C₁ to C₁₀ alkyl or C₂ to C₁₀ alkenyl and alkynyl. Particularly preferred are O[(CH₂)_nO]_mCH₃, O(CH₂)_nOCH₃, O(CH₂)_nNH₂, O(CH₂)_nCH₃, O(CH₂)_nONH₂, and O(CH₂)_nON[(CH₂)_nCH₃]₂, where n and m are from 1 to about 10. Other preferred oligonucleotides comprise one of the
25 following at the 2' position: C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA
30 cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred
35 modification includes 2'-methoxyethoxy (2'-O-CH₂CH₂OCH₃, also

-15-

known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., *Helv. Chim. Acta*, 1995, 78, 486-504) i.e., an alkoxyalkoxy group. A further preferred modification includes 2'-dimethylaminooxyethoxy, i.e., a $O(CH_2)_2ON(CH_3)_2$ group, also
5 known as 2'-DMAOE, as described in United States patent application Serial Number 09/016,520, filed on January 30, 1998, which is commonly owned with the instant application and the contents of which are herein incorporated by reference.

10 Other preferred modifications include 2'-methoxy (2'-O-CH₃), 2'-aminopropoxy (2'-OCH₂CH₂CH₂NH₂) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in
15 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotides. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar
20 structures include, but are not limited to, U.S.:
4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878;
5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811;
5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053;
5,639,673; 5,646,265; 5,658,873; 5,670,633; and 5,700,920,
25 each of which is incorporated herein by reference.

Oligonucleotides may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and
30 guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-Me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of
35 adenine and guanine, 2-propyl and other alkyl derivatives

-16-

of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further nucleobases include those disclosed in U.S. Patent 3,687,808, those disclosed in *The Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., *Angewandte Chemie*, International Edition, 1991, 30, 613, and those disclosed by Sanghvi, Y.S., Crooke, S.T. and Lebleu, B. eds., *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 289-302. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and Lebleu, B., eds., *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

Representative United States patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, U.S.: 3,687,808; 4,845,205;

-17-

5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272;
5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711;
5,552,540; 5,587,469; 5,594,121; 5,596,091; 5,614,617;
5,681,941; and 5,750,692, each of which is herein

5 incorporated by reference.

Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates which enhance the activity, cellular distribution or cellular
10 uptake of the oligonucleotide. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, 1989, 86, 6553-6556), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1994, 4, 1053-1060), a thioether, e.g., hexyl-
15 S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, 1992, 660, 306-309; Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1993, 3, 2765-2770), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, 1992, 20, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-
20 Behmoaras et al., *EMBO J.*, 1991, 10, 1111-1118; Kabanov et al., *FEBS Lett.*, 1990, 259, 327-330; Svinarchuk et al., *Biochimie*, 1993, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al.,
25 *Tetrahedron Lett.*, 1995, 36, 3651-3654; Shea et al., *Nucl. Acids Res.*, 1990, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., *Nucleosides & Nucleotides*, 1995, 14, 969-973), or adamantane acetic acid (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651-3654),
30 a palmityl moiety (Mishra et al., *Biochim. Biophys. Acta*, 1995, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl-oxysterol moiety (Crooke et al., *J. Pharmacol. Exp. Ther.*, 1996, 277, 923-937).

-18-

Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but are not limited to, U.S.: 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717, 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941, each of which is herein incorporated by reference.

It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single compound or even at a single nucleoside within an oligonucleotide. The present invention also includes antisense compounds which are chimeric compounds. "Chimeric" antisense compounds or "chimeras," in the context of this invention, are antisense compounds, particularly oligonucleotides, which contain two or more chemically distinct regions, each made up of at least one monomer unit *i.e.*, a nucleotide in the case of an oligonucleotide compound. These oligonucleotides typically contain at least one region wherein the oligonucleotide is modified so as to confer upon the oligonucleotide increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. An additional region of the oligonucleotide may serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is

-19-

a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of oligonucleotide inhibition of gene expression. Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

Chimeric antisense compounds of the invention may be formed as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics as described above. Such compounds have also been referred to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures include, but are not limited to, U.S.: 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922, each of which is herein incorporated by reference.

The antisense compounds used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, CA). Any other means for such synthesis known in the art may additionally or alternatively be employed. It is well known to use similar techniques to prepare oligonucleotides such as the phosphorothioates and alkylated derivatives.

The antisense compounds of the invention are synthesized *in vitro* and do not include antisense

-20-

compositions of biological origin, or genetic vector constructs designed to direct the *in vivo* synthesis of antisense molecules.

The compounds of the invention may also be admixed, 5 encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as for example, liposomes, receptor targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative 10 United States patents that teach the preparation of such uptake, distribution and/or absorption assisting formulations include, but are not limited to, U.S.: 5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291; 5,543,158; 5,547,932; 5,583,020; 5,591,721; 4,426,330; 15 4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170; 5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978; 5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259; 5,543,152; 5,556,948; 5,580,575; and 5,595,756, each of which is herein incorporated by reference.

20 The antisense compounds of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal including a human, is capable of providing (directly or indirectly) the biologically active metabolite 25 or residue thereof. Accordingly, for example, the disclosure is also drawn to prodrugs and pharmaceutically acceptable salts of the compounds of the invention, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents.

30 The term "prodrug" indicates a therapeutic agent that is prepared in an inactive form that is converted to an active form (*i.e.*, drug) within the body or cells thereof by the action of endogenous enzymes or other chemicals and/or conditions. In particular, prodrug versions of the 35 oligonucleotides of the invention are prepared as CATE

-21-

[(S-acetyl-2-thioethyl) phosphatel derivatives according to the methods disclosed in WO 93/24510 or in WO 94/26764.

The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto (see, for example, Berge et al., "Pharmaceutical Salts," *J. of Pharma Sci.*, 1977, 66, 1-19).

For oligonucleotides, preferred examples of pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed from elemental anions such as chlorine, bromine, and iodine.

The antisense compounds of the present invention can be utilized for diagnostics, therapeutics, prophylaxis and as research reagents and kits. For therapeutics, an animal, preferably a human, suspected of having a disease or disorder which can be treated by modulating the expression of one or more members of the TRAF family is treated by administering antisense compounds in accordance with this invention. The compounds of the invention can be utilized in pharmaceutical compositions by adding an

-22-

effective amount of an antisense compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the antisense compounds and methods of the invention may also be useful prophylactically, e.g., to prevent or delay

5 infection, inflammation or tumor formation, for example.

The antisense compounds of the invention are useful for research and diagnostics, because these compounds hybridize to nucleic acids encoding one or more selected TRAFs, enabling sandwich and other assays to easily be

10 constructed to exploit this fact. Hybridization of the antisense oligonucleotides of the invention with a nucleic acid encoding one or more TRAFs can be detected by means known in the art. Such means may include conjugation of an enzyme to the oligonucleotide, radiolabelling of the

15 oligonucleotide or any other suitable detection means. Kits using such detection means for detecting the level of TRAF in a sample may also be prepared.

The present invention also includes pharmaceutical compositions and formulations which include the antisense

20 compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including

25 ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; (intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral

30 administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be

35 particularly useful for oral administration.

-23-

Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful.

Compositions and formulations for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, capsules, sachets or tablets. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable.

Compositions and formulations for parenteral, intrathecal or intraventricular administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients.

For example, pharmaceutical compositions and/or formulations comprising the oligonucleotides of the present invention may include penetration enhancers in order to enhance the alimentary delivery of the oligonucleotides. Penetration enhancers may be classified as belonging to one of five broad categories, i.e., fatty acids, bile salts, chelating agents, surfactants and non-surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, 8, 91-192; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33). One or more penetration enhancers from one or more of these broad categories may be included.

Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprato, tricaprato, recinleate, monoolein (a.k.a. 1-monooleoyl-rac-

-24-

glycerol), dilaurin, caprylic acid, arachidonic acid, glyceryl 1-monocaprate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, mono- and di-glycerides and physiologically acceptable salts thereof (i.e., oleate, 5 laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, 8, 2, 91-192; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1, 1-33; El-Hariri et al., *J. Pharm.* 10 *Pharmacol.*, 1992, 44, 651-654). Examples of some presently preferred fatty acids are sodium caprate and sodium laurate, used singly or in combination at concentrations of 0.5 to 5%.

The physiological roles of bile include the 15 facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 In: *Goodman & Gilman's The Pharmacological Basis of Therapeutics*, 9th Ed., Hardman et al., eds., McGraw-Hill, New York, NY, 1996, pages 934-935). Various natural bile salts, and their 20 synthetic derivatives, act as penetration enhancers. Thus, the term "bile salt" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives.

Regardless of the method by which the antisense 25 compounds of the invention are introduced into a patient, colloidal dispersion systems may be used as delivery vehicles to enhance the *in vivo* stability of the compounds and/or to target the compounds to a particular organ, tissue or cell type. Colloidal dispersion systems include, 30 but are not limited to, macromolecule complexes, nanocapsules, microspheres, beads and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, liposomes and lipid:oligonucleotide complexes of uncharacterized structure. A preferred colloidal

-25-

dispersion system is a plurality of liposomes. Liposomes are microscopic spheres having an aqueous core surrounded by one or more outer layer(s) made up of lipids arranged in a bilayer configuration (see, generally, Chonn et al.,
5 *Current Op. Biotech.*, 1995, 6, 698-708).

Liposome preparation is described in pending United States patent application 08/961,469, filed on October 31, 1997, which is commonly owned with the instant application and which is herein incorporated by reference.

10 Certain embodiments of the invention provide for liposomes and other compositions containing (a) one or more antisense compounds and (b) one or more other chemotherapeutic agents which function by a non-antisense mechanism. Other non-antisense chemotherapeutic agents are
15 also within the scope of this invention. Two or more combined compounds may be used together or sequentially.

In another related embodiment, compositions of the invention may contain one or more antisense compounds, particularly oligonucleotides, targeted to a first nucleic
20 acid and one or more additional antisense compounds targeted to a second nucleic acid target. Two or more combined compounds may be used together or sequentially.

The formulation of therapeutic compositions and their subsequent administration is believed to be within the
25 skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules
30 can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and
35 can generally be estimated based on EC₅₀s found to be

-26-

effective in *in vitro* and *in vivo* animal models. In general, dosage is from 0.01 μg to 100 g per kg of body weight, and may be given once or more daily, weekly, monthly or yearly, or even once every 2 to 20 years.

- 5 Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy
10 to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses, ranging from 0.01 μg to 100 g per kg of body weight, once or more daily, to once every 20 years.

While the present invention has been described with
15 specificity in accordance with certain of its preferred embodiments, the following examples serve only to illustrate the invention and are not intended to limit the same.

EXAMPLES

20 Example 1

Nucleoside Phosphoramidites for Oligonucleotide Synthesis
Deoxy and 2'-alkoxy amidites

- 2'-Deoxy and 2'-methoxy beta-cyanoethyl-diisopropyl phosphoramidites are purchased from commercial sources
25 (e.g., Chemgenes, Needham, MA or Glen Research, Inc. Sterling VA). Other 2'-O-alkoxy substituted nucleoside amidites are prepared as described in U.S. Patent 5,506,351, herein incorporated by reference. For oligonucleotides synthesized using 2'-alkoxy amidites, the
30 standard cycle for unmodified oligonucleotides is utilized, except the wait step after pulse delivery of tetrazole and base is increased to 360 seconds.

Oligonucleotides containing 5-methyl-2'-deoxycytidine (5-Me-C) nucleotides were synthesized according to

-27-

published methods (Sanghvi, et. al., *Nucleic Acids Research*, 1993, 21, 3197-3203] using commercially available phosphoramidites (Glen Research, Sterling VA or ChemGenes, Needham, MA).

5 **2'-Fluoro amidites**

2'-Fluorodeoxyadenosine amidites

 2'-fluoro oligonucleotides are synthesized as described previously by Kawasaki, et. al., *J. Med. Chem.*, 1993, 36, 831-841 and U.S. Patent 5,670,633, herein
10 incorporated by reference. Briefly, the protected nucleoside N6-benzoyl-2'-deoxy-2'-fluoroadenosine are synthesized utilizing commercially available 9-beta-D-arabinofuranosyladenine as starting material and by
15 modifying literature procedures whereby the 2'-alpha-fluoro atom is introduced by a S_N2-displacement of a 2'-beta-trityl group. Thus N6-benzoyl-9-beta-D-arabinofuranosyladenine is selectively protected in moderate yield as the 3',5'-ditetrahydropyranyl (THP) intermediate. Deprotection of
20 the THP and N6-benzoyl groups is accomplished using standard methodologies and standard methods are used to obtain the 5'-dimethoxytrityl-(DMT) and 5'-DMT-3'-phosphoramidite intermediates.

2'-Fluorodeoxyguanosine

 The synthesis of 2'-deoxy-2'-fluoroguanosine is
25 accomplished using tetraisopropylidisiloxanyl (TPDS) protected 9-beta-D-arabinofuranosylguanine as starting material, and conversion to the intermediate diisobutyryl-arabinofuranosylguanosine. Deprotection of the TPDS group is followed by protection of the hydroxyl group with THP to
30 give diisobutyryl di-THP protected arabinofuranosylguanine. Selective O-deacylation and triflation is followed by treatment of the crude product with fluoride, then deprotection of the THP groups. Standard methodologies are used to obtain the 5'-DMT- and 5'-DMT-3'-phosphoramidites.

2'-Fluorouridine

Synthesis of 2'-deoxy-2'-fluorouridine is accomplished by the modification of a literature procedure in which
5 2,2'-anhydro-1-beta-D-arabinofuranosyluracil is treated with 70% hydrogen fluoride-pyridine. Standard procedures are used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

2'-Fluorodeoxycytidine

10 2'-deoxy-2'-fluorocytidine is synthesized via amination of 2'-deoxy-2'-fluorouridine, followed by selective protection to give N4-benzoyl-2'-deoxy-2'-fluorocytidine. Standard procedures are used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

15 2'-O-(2-Methoxyethyl) modified amidites

2'-O-Methoxyethyl-substituted nucleoside amidites were prepared as follows, or alternatively, as per the methods of Martin, P., *Helvetica Chimica Acta*, 1995, 78, 486-504.

20 **2,2'-Anhydro[1-(beta-D-arabinofuranosyl)-5-methyluridine]**

5-Methyluridine (ribosylthymine, commercially available through Yamasa, Choshi, Japan) (72.0 g, 0.279 M), diphenylcarbonate (90.0 g, 0.420 M) and sodium bicarbonate (2.0 g, 0.024 M) were added to DMF (300 mL). The mixture
25 was heated to reflux, with stirring, allowing the evolved carbon dioxide gas to be released in a controlled manner. After 1 hour, the slightly darkened solution was concentrated under reduced pressure. The resulting syrup was poured into diethylether (2.5 L), with stirring. The
30 product formed a gum. The ether was decanted and the residue was dissolved in a minimum amount of methanol (ca. 400 mL). The solution was poured into fresh ether (2.5 L) to yield a stiff gum. The ether was decanted and the gum dried in a vacuum oven (60°C at 1 mm Hg for 24 hours) to

-29-

give a solid that was crushed to a light tan powder (57 g, 85% crude yield). The NMR spectrum was consistent with the structure, contaminated with phenol as its sodium salt (ca. 5%). The material was used as is for further reactions (or
5 purified further by column chromatography using a gradient of methanol in ethyl acetate (10-25%) to give a white solid, mp 222-4°C).

2'-O-Methoxyethyl-5-methyluridine

2.2'-Anhydro-5-methyluridine (195 g, 0.81 M), tris(2-
10 methoxyethyl)borate (231 g, 0.98 M) and 2-methoxyethanol (1.2 L) were added to a 2 L stainless steel pressure vessel and placed in a pre-heated oil bath at 160°C. After heating for 48 hours at 155-160°C, the vessel was opened and the solution evaporated to dryness and triturated with MeOH
15 (200 mL). The residue was suspended in hot acetone (1 L). The insoluble salts were filtered, washed with acetone (150 mL) and the filtrate evaporated. The residue (280 g) was dissolved in CH₃CN (600 mL) and evaporated. A silica gel column (3 kg) was packed in CH₂Cl₂/Acetone/MeOH (20:5:3)
20 containing 0.5% Et₃NH. The residue was dissolved in CH₂Cl₂ (250 mL) and adsorbed onto silica (150 g) prior to loading onto the column. The product was eluted with the packing solvent to give 160 g (63%) of product. Additional material was obtained by reworking impure fractions.

25 **2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine**

2'-O-Methoxyethyl-5-methyluridine (160 g, 0.506 M) was co-evaporated with pyridine (250 mL) and the dried residue dissolved in pyridine (1.3 L). A first aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and
30 the mixture stirred at room temperature for one hour. A second aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the reaction stirred for an additional one hour. Methanol (170 mL) was then added to stop the reaction. HPLC showed the presence of approximately 70%
35 product. The solvent was evaporated and triturated with

-30-

CH₃CN (200 mL). The residue was dissolved in CHCl₃ (1.5 L) and extracted with 2x500 mL of saturated NaHCO₃ and 2x500 mL of saturated NaCl. The organic phase was dried over Na₂SO₄, filtered and evaporated. The residue was purified on a 3.5 kg silica gel column, packed and eluted with EtOAc/Hexane/Acetone (5:5:1) containing 0.5% Et₃NH. The pure fractions were evaporated to give 164 g of product. Approximately 20 g additional was obtained from the impure fractions to give a total yield of 183 g (57%).

10 3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (106 g, 0.167 M), DMF/pyridine (750 mL of a 3:1 mixture prepared from 562 mL of DMF and 188 mL of pyridine) and acetic anhydride (24.38 mL, 0.258 M) were combined and stirred at room temperature for 24 hours. The reaction was monitored by tlc by first quenching the tlc sample with the addition of MeOH. Upon completion of the reaction, as judged by tlc, MeOH (50 mL) was added and the mixture evaporated at 35°C. The residue was dissolved in CHCl₃ (800 mL) and extracted with 2x200 mL of saturated sodium bicarbonate and 2x200 mL of saturated NaCl. The water layers were back extracted with 200 mL of CHCl₃. The combined organics were dried with sodium sulfate and evaporated to give 122 g of residue (approx. 90% product). The residue was purified on a 3.5 kg silica gel column and eluted using EtOAc/Hexane(4:1). Pure product fractions were evaporated to yield 96 g (84%). An additional 1.5 g was recovered from later fractions.

25 3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine

A first solution was prepared by dissolving 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (96 g, 0.144 M) in CH₃CN (700 mL) and set aside. Triethylamine (189 mL, 1.44 M) was added to a

-31-

solution of triazole (90 g, 1.3 M) in CH_3CN (1 L), cooled to -5°C and stirred for 0.5 hours using an overhead stirrer. POCl_3 was added dropwise, over a 30 minute period, to the stirred solution maintained at $0-10^\circ\text{C}$, and the resulting mixture stirred for an additional 2 hours. The first solution was added dropwise, over a 45 minute period, to the latter solution. The resulting reaction mixture was stored overnight in a cold room. Salts were filtered from the reaction mixture and the solution evaporated. The residue was dissolved in EtOAc (1 L) and the insoluble solids removed by filtration. The filtrate was washed with 1x300 mL of NaHCO_3 and 2x300 mL of saturated NaCl, dried over sodium sulfate and evaporated. The residue was triturated with EtOAc to give the title compound.

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine

A solution of 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine (103 g, 0.141 M) in dioxane (500 mL) and NH_4OH (30 mL) was stirred at room temperature for 2 hours. The dioxane solution was evaporated and the residue azeotroped with MeOH (2x200 mL). The residue was dissolved in MeOH (300 mL) and transferred to a 2 liter stainless steel pressure vessel. MeOH (400 mL) saturated with NH_3 gas was added and the vessel heated to 100°C for 2 hours (tlc showed complete conversion). The vessel contents were evaporated to dryness and the residue was dissolved in EtOAc (500 mL) and washed once with saturated NaCl (200 mL). The organics were dried over sodium sulfate and the solvent evaporated to give 85 g (95%) of the title compound.

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (85 g, 0.134 M) was dissolved in DMF (800 mL) and benzoic anhydride (37.2 g, 0.165 M) is added with stirring.

-32-

The mixture was stirred for 3 hours (tlc showing the reaction to be approximately 95% complete). The solvent was evaporated and the residue azeotroped with MeOH (200 mL). The residue was then dissolved in CHCl₃ (700 mL) and
5 extracted with saturated NaHCO₃ (2x300 mL) and saturated NaCl (2x300 mL), dried over MgSO₄, and evaporated to give a residue (96 g). The residue was chromatographed on a 1.5 kg silica column using EtOAc/Hexane (1:1) containing 0.5% Et₃NH as the eluting solvent. The pure product fractions
10 were evaporated to give 90 g (90%) of the title compound.

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine-3'-amidite

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (74 g, 0.10 M) was dissolved in CH₂Cl₂ (1 L).
15 Tetrazole diisopropylamine (7.1 g) and 2-cyanoethoxy-tetra-(isopropyl)phosphite (40.5 mL, 0.123 M) were added with stirring, under a nitrogen atmosphere. The resulting mixture was stirred for 20 hours at room temperature (tlc showing the reaction to be 95% complete). The reaction
20 mixture was extracted with saturated NaHCO₃ (1x300 mL) and saturated NaCl (3x300 mL). The aqueous washes were back-extracted with CH₂Cl₂ (300 mL), and the extracts were combined, dried over MgSO₄, and concentrated. The residue
25 obtained was chromatographed on a 1.5 kg silica column using EtOAc/Hexane (3:1) as the eluting solvent. The pure fractions were combined to give 90.6 g (87%) of the title compound.

2'-(Aminooxyethyl) nucleoside amidites and 2'-(dimethylaminooxyethyl) nucleoside amidites

30 Aminooxyethyl and dimethylaminooxyethyl amidites are prepared as per the methods described in United States patent applications serial number 10/037,143, filed February 14, 1998, and serial number 09/016,520, filed January 30, 1998, each of which is herein incorporated by
35 reference.

-33-

Example 2**Oligonucleotide synthesis**

Unsubstituted and substituted phosphodiester (P=O) oligonucleotides are synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) using standard phosphoramidite chemistry with oxidation by iodine.

Phosphorothioates (P=S) were synthesized as for the phosphodiester oligonucleotides except the standard oxidation bottle was replaced by 0.2 M solution of 3H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the stepwise thiation of the phosphite linkages. The thiation wait step was increased to 68 seconds and was followed by the capping step. After cleavage from the CPG column and deblocking in concentrated ammonium hydroxide at 55°C (18 hours), the oligonucleotides were purified by precipitating twice with 2.5 volumes of ethanol from a 0.5 M NaCl solution.

Phosphinate oligonucleotides are prepared as described in U.S. Patent 5,508,270, herein incorporated by reference.

Alkyl phosphonate oligonucleotides are prepared as described in U.S. Patent 4,469,863, herein incorporated by reference.

3'-Deoxy-3'-methylene phosphonate oligonucleotides are prepared as described in U.S. Patents 5,610,289 or 5,625,050, herein incorporated by reference.

Phosphoramidite oligonucleotides are prepared as described in U.S. Patent, 5,256,775 or U.S. Patent 5,366,878, herein incorporated by reference.

Alkylphosphonothioate oligonucleotides are prepared as described in published PCT applications PCT/US94/00902 and PCT/US93/06976 (published as WO 94/17093 and WO 94/02499, respectively), herein incorporated by reference.

3'-Deoxy-3'-amino phosphoramidate oligonucleotides are prepared as described in U.S. Patent 5,476,925, herein incorporated by reference.

-34-

Phosphotriester oligonucleotides are prepared as described in U.S. Patent 5,023,243, herein incorporated by reference.

Borano phosphate oligonucleotides are prepared as
5 described in U.S. Patents 5,130,302 and 5,177,198, herein incorporated by reference.

Example 3

Oligonucleoside Synthesis

Methylenemethylimino linked oligonucleosides, also
10 identified as MMI linked oligonucleosides, methylenedi-
methylhydrazo linked oligonucleosides, also identified as
MDH linked oligonucleosides, methylenecarbonylamino linked
oligonucleosides, also identified as amide-3 linked
oligonucleosides, and methyleneaminocarbonyl linked oligo-
15 nucleosides, also identified as amide-4 linked oligonucleo-
sides, as well as mixed backbone compounds having, for
instance, alternating MMI and P=O or P=S linkages are
prepared as described in U.S. Patents 5,378,825, 5,386,023,
5,489,677, 5,602,240 and 5,610,289, all of which are
20 herein incorporated by reference.

Formacetal and thioformacetal linked oligonucleosides
are prepared as described in U.S. Patents 5,264,562 and
5,264,564, herein incorporated by reference.

Ethylene oxide linked oligonucleosides are prepared as
25 described in U.S. Patent 5,223,618, herein incorporated by
reference.

Example 4

PNA Synthesis

Peptide nucleic acids (PNAs) are prepared in
30 accordance with any of the various procedures referred to
in *Peptide Nucleic Acids (PNA): Synthesis, Properties and
Potential Applications, Bioorganic & Medicinal Chemistry,*
1996, 4, 5-23. They may also be prepared in accordance

-35-

with U.S. Patents 5,539,082, 5,700,922, and 5,719,262, herein incorporated by reference.

Example 5**Synthesis of Chimeric Oligonucleotides**

5 Chimeric oligonucleotides, oligonucleosides or mixed oligonucleotides/oligonucleosides of the invention can be of several different types. These include a first type wherein the "gap" segment of linked nucleosides is positioned between 5' and 3' "wing" segments of linked
10 nucleosides and a second "open end" type wherein the "gap" segment is located at either the 3' or the 5' terminus of the oligomeric compound. Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of the second type are
15 also known in the art as "hemimers" or "wingmers".

[2'-O-Me]--[2'-deoxy]--[2'-O-Me] Chimeric

Phosphorothioate Oligonucleotides

Chimeric oligonucleotides having 2'-O-alkyl phosphorothioate and 2'-deoxy phosphorothioate oligo-
20 nucleotide segments are synthesized using an Applied Biosystems automated DNA synthesizer Model 380B, as above. Oligonucleotides are synthesized using the automated synthesizer and 2'-deoxy-5'-dimethoxytrityl-3'-O-phosphoramidite for the DNA portion and 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidite for 5' and 3' wings. The
25 standard synthesis cycle is modified by increasing the wait step after the delivery of tetrazole and base to 600 seconds repeated four times for RNA and twice for 2'-O-methyl. The fully protected oligonucleotide is cleaved
30 from the support and the phosphate group is deprotected in 3:1 Ammonia/Ethanol at room temperature overnight then lyophilized to dryness. Treatment in methanolic ammonia for 24 hours at room temperature is then done to deprotect all bases and sample was again lyophilized to dryness. The
35 pellet is resuspended in 1M TBAF in THF for 24 hours at

-36-

room temperature to deprotect the 2' positions. The reaction is then quenched with 1M TEAA and the sample is then reduced to 1/2 volume by rotovac before being desalted on a G25 size exclusion column. The oligo recovered is then analyzed spectrophotometrically for yield and for purity by capillary electrophoresis and by mass spectrometry.

- [2'-O-(2-Methoxyethyl)]--[2'-deoxy]--[2'-O-(Methoxyethyl)] Chimeric Phosphorothioate Oligonucleotides
- [2'-O-(2-methoxyethyl)]--[2'-deoxy]--[2'-O-(methoxyethyl)] chimeric phosphorothioate oligonucleotides were prepared as per the procedure above for the 2'-O-methyl chimeric oligonucleotide, with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites.
- [2'-O-(2-Methoxyethyl)Phosphodiester]--[2'-deoxy Phosphorothioate]--[2'-O-(2-Methoxyethyl) Phosphodiester] Chimeric Oligonucleotides
- [2'-O-(2-methoxyethyl phosphodiester)]--[2'-deoxy phosphorothioate]--[2'-O-(methoxyethyl) phosphodiester] chimeric oligonucleotides were prepared as per the above procedure for the 2'-O-methyl chimeric oligonucleotide with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites, oxidization with iodine to generate the phosphodiester internucleotide linkages within the wing portions of the chimeric structures and sulfurization utilizing 3,4-dithiolane-2-one 1,1 dioxides (DeaCage Reagent) to generate the phosphorothioate internucleotide linkages for the center gap.
- Other chimeric oligonucleotides, chimeric oligonucleosides and mixed chimeric oligonucleotides/oligonucleosides are synthesized according to United States patent 5,623,065, herein incorporated by reference.

-37-

Example 6**Oligonucleotide Isolation**

After cleavage from the controlled pore glass column (Applied Biosystems) and deblocking in concentrated ammonium hydroxide at 55°C for 18 hours, the oligonucleotides or oligonucleosides were purified by precipitation twice out of 0.5 M NaCl with 2.5 volumes ethanol. Synthesized oligonucleotides were analyzed by polyacrylamide gel electrophoresis on denaturing gels and judged to be at least 85% full length material. The relative amounts of phosphorothioate and phosphodiester linkages obtained in synthesis were periodically checked by ³¹P nuclear magnetic resonance spectroscopy, and for some studies oligonucleotides are purified by HPLC, as described by Chiang et al., *J. Biol. Chem.* 1991, 266, 18162-18171. Results obtained with HPLC-purified material are similar to those obtained with non-HPLC purified material.

Example 7**Oligonucleotide Synthesis - 96 Well Plate Format**

Oligonucleotides were synthesized via solid phase P(III) phosphoramidite chemistry on an automated synthesizer capable of assembling 96 sequences simultaneously in a standard 96 well format. Phosphodiester internucleotide linkages were afforded by oxidation with aqueous iodine. Phosphorothioate internucleotide linkages were generated by sulfurization utilizing 3,4-dihydro-2H-benzodithiole-3-one 1,1 dioxide (Beaucage Reagent) in anhydrous acetonitrile. Standard base-protected beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial vendors (e.g. PE-Applied Biosystems, Foster City, CA, or Pharmacia, Piscataway, NJ). Non-standard nucleosides were synthesized as per known literature or patented methods. They were utilized as base protected beta-cyanoethyl-diisopropyl phosphoramidites.

-38-

Oligonucleotides were cleaved from support and deprotected with concentrated NH_4OH at elevated temperature (55-60°C) for 12-16 hours and the released product then dried *in vacuo*. The dried product was then re-suspended in
5 sterile water to afford a master plate from which all analytical and test plate samples were then diluted utilizing robotic pipettors.

Example 8**Oligonucleotide Analysis - 96 Well Plate Format**

10 The concentration of oligonucleotide in each well was assessed by dilution of samples and UV absorption spectroscopy. The full-length integrity of the individual products was evaluated by capillary electrophoresis (CE) in either the 96 well format (Beckman P/ACE™ MDQ) or, for
15 individually prepared samples, on a commercial CE apparatus (e.g., Beckman P/ACE™ 5000, ABI 270). Base and backbone composition was confirmed by mass analysis of the compounds utilizing electrospray-mass spectroscopy. All assay test
20 plates were diluted from the master plate using single and multi-channel robotic pipettors. Plates were judged to be acceptable if at least 85% of the compounds on the plate were at least 85% full length.

Example 9**Cell culture and oligonucleotide treatment**

25 The effect of antisense compounds on target nucleic acid expression can be tested in any of a variety of cell types provided that the target nucleic acid is present at measurable levels. This can be routinely determined using, for example, PCR, RNase protection assay (RPA) or Northern
30 blot analysis. The following four cell types are provided for illustrative purposes, but other cell types can be routinely used.

-39-

T-24 cells:

The transitional cell bladder carcinoma cell line T-24 is obtained from the American Type Culture Collection (ATCC) (Manassas, VA). T-24 cells are routinely cultured in complete McCoy's 5A basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells are routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells are seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

For Northern blotting or other analysis, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

A549 cells:

The human lung carcinoma cell line A549 is obtained from the American Type Culture Collection (ATCC) (Manassas, VA). A549 cells are routinely cultured in DMEM basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells are routinely passaged by trypsinization and dilution when they reach 90% confluence.

NHDF cells:

Human neonatal dermal fibroblast (NHDF) are obtained from the Clonetics Corporation (Walkersville MD). NHDFs are routinely maintained in Fibroblast Growth Medium (Clonetics Corporation, Walkersville, MD) supplemented as

-40-

recommended by the supplier. Cells are maintained for up to 10 passages as recommended by the supplier.

HEK cells:

5 Human embryonic keratinocytes (HEK) are obtained from the Clonetics Corporation (Walkersville, MD). HEKs are routinely maintained in Keratinocyte Growth Medium (Clonetics Corporation, Walkersville, MD) formulated as recommended by the supplier. Cells are routinely
10 maintained for up to 10 passages as recommended by the supplier.

Treatment with antisense compounds:

When cells reach 80% confluency, they are treated with
15 oligonucleotide. For cells grown in 96-well plates, wells are washed once with 200 μ L OPTI-MEM™-1 reduced-serum medium (Gibco BRL) and then treated with 130 μ L of OPTI-MEM™-1 containing 3.75 μ g/mL LIPOFECTIN™ (Gibco BRL) and the desired oligonucleotide at a final concentration of 150
20 nM. After 4 hours of treatment, the medium is replaced with fresh medium. Cells are harvested 16 hours after oligonucleotide treatment.

Example 10

Analysis of oligonucleotide inhibition of TRAF expression

25 Antisense modulation of TRAF expression can be assayed in a variety of ways known in the art. For example, TRAF mRNA levels can be quantitated by, e.g., Northern blot analysis, RNase protection assay (RPA), competitive polymerase chain reaction (PCR), or real-time PCR (RT-PCR).
30 RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. Methods of RNA isolation are taught in, for example, Ausubel, et al., *Current Protocols in Molecular Biology*, Volume 1, John Wiley & Sons, Inc., 1993, pp. 4.1.1-4.2.9 and 4.5.1-4.5.3. Northern blot analysis is
35 routine in the art and is taught in, for example, Ausubel,

-41-

et al., *Current Protocols in Molecular Biology*, Volume 1, John Wiley & Sons, Inc., 1996, pp. 4.2.1-4.2.9. Real-time quantitative (PCR) can be conveniently accomplished using the commercially available ABI PRISM™ 7700 Sequence
5 Detection System, available from PE-Applied Biosystems, Foster City, CA and used according to manufacturer's instructions. Other methods of PCR are also known in the art.

TRAF protein levels can be quantitated in a variety of
10 ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), ELISA or fluorescence-activated cell sorting (FACS). Antibodies directed to TRAF can be identified and obtained from a variety of sources, such as the MSRS catalog of antibodies
15 (Aerie Corporation, Birmingham, MI), or can be prepared via conventional antibody generation methods. Methods for preparation of polyclonal antisera are taught in, for example, Ausubel, et al., *Current Protocols in Molecular Biology*, Volume 2, John Wiley & Sons, Inc., 1997, pp.
20 11.12.1-11.12.9. Preparation of monoclonal antibodies is taught in, for example, Ausubel, et al., *Current Protocols in Molecular Biology*, Volume 2, John Wiley & Sons, Inc., 1997, pp. 11.4.1-11.11.5.

Immunoprecipitation methods are standard in the art
25 and can be found at, for example, Ausubel, et al., *Current Protocols in Molecular Biology*, Volume 2, John Wiley & Sons, Inc., 1998, pp. 11.4.1-11.11.5. Western blot (immunoblot) analysis is standard in the art and can be found at, for example, Ausubel, et al., *Current Protocols in Molecular Biology*, Volume 2, John Wiley & Sons, Inc.,
30 1997, pp. 10.8.1-10.8.21. Enzyme-linked immunosorbent assays (ELISA) are standard in the art and can be found at, for example, Ausubel, et al., *Current Protocols in*

-42-

Molecular Biology, Volume 2, John Wiley & Sons, Inc., 1991, pp. 11.2.1-11.2.22.

Example 11**Poly(A)+ mRNA isolation**

- 5 Poly(A)+ mRNA is isolated according to Miura et al., *Clin. Chem.*, 1996, 42, 1758-1764. Other methods for poly(A)+ mRNA isolation are taught in, for example, Ausubel, et al., *Current Protocols in Molecular Biology*, Volume 1, John Wiley & Sons, Inc., 1993, pp. 4.5.1-4.5.3.
- 10 Briefly, for cells grown on 96-well plates, growth medium is removed from the cells and each well is washed with 200 μ L cold PBS. 60 μ L lysis buffer (10 mM Tris-HCl, pH 7.6, 1 mM EDTA, 0.5 M NaCl, 0.5% NP-40, 20 mM vanadyl-ribonucleoside complex) is added to each well, the plate is
- 15 gently agitated and then incubated at room temperature for 5 minutes. 55 μ L of lysate is transferred to Oligo d(T) coated 96-well plates (AGCT Inc., Irvine, CA). Plates are incubated for 60 minutes at room temperature, washed 3 times with 200 μ L of wash buffer (10 mM Tris-HCl pH 7.6, 1 mM EDTA, 0.3 M NaCl). After the final wash, the plate is
- 20 blotted on paper towels to remove excess wash buffer and then air-dried for 5 minutes. 60 μ L of elution buffer (5 mM Tris-HCl pH 7.6), preheated to 70°C is added to each well, the plate is incubated on a 90°C hot plate for 5 minutes,
- 25 and the eluate is then transferred to a fresh 96-well plate.

Cells grown on 100 mm or other standard plates may be treated similarly, using appropriate volumes of all solutions.

30 Example 12**Total RNA Isolation**

Total mRNA was isolated using an RNEASY 96™ kit and buffers purchased from Qiagen Inc. (Valencia, CA) following the manufacturer's recommended procedures. Briefly, for

-43-

cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200 μ L cold PBS. 100 μ L Buffer RLT was added to each well and the plate vigorously agitated for 20 seconds. 100 μ L of 70% ethanol was then added to each well and the contents mixed by pipetting three times up and down. The samples were then transferred to the RNEASY 96™ well plate attached to a QIAVAC™ manifold fitted with a waste collection tray and attached to a vacuum source. Vacuum was applied for 15 seconds. 1 mL of Buffer RW1 was added to each well of the RNEASY 96™ plate and the vacuum again applied for 15 seconds. 1 mL of Buffer RPE was then added to each well of the RNEASY 96™ plate and the vacuum applied for a period of 15 seconds. The Buffer RPE wash was then repeated and the vacuum was applied for an additional 10 minutes. The plate was then removed from the QIAVAC™ manifold and blotted dry on paper towels. The plate was then re-attached to the QIAVAC™ manifold fitted with a collection tube rack containing 1.2 mL collection tubes. RNA was then eluted by pipetting 60 μ L water into each well, incubating 1 minute, and then applying the vacuum for 30 seconds. The elution step was repeated with an additional 60 μ L water.

Example 13**Real-time Quantitative PCR Analysis of TRAF mRNA Levels**

Quantitation of TRAF mRNA levels is determined by real-time quantitative PCR using the ABI PRISM™ 7700 Sequence Detection System (PE-Applied Biosystems, Foster City, CA) according to manufacturer's instructions. This is a closed-tube, non-gel-based, fluorescence detection system which allows high-throughput quantitation of polymerase chain reaction (PCR) products in real-time. As opposed to standard PCR, in which amplification products are quantitated after the PCR is completed, products in real-time quantitative PCR are quantitated as they accumulate. This is accomplished by including in the PCR

-44-

reaction an oligonucleotide probe that anneals specifically between the forward and reverse PCR primers, and contains two fluorescent dyes. A reporter dye (e.g., JOE or FAM, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 5' end of the probe and a quencher dye (e.g., TAMRA, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 3' end of the probe. When the probe and dyes are intact, reporter dye emission is quenched by the proximity of the 3' quencher dye. During amplification, annealing of the probe to the target sequence creates a substrate that can be cleaved by the 5'-exonuclease activity of Taq polymerase. During the extension phase of the PCR amplification cycle, cleavage of the probe by Taq polymerase releases the reporter dye from the remainder of the probe (and hence from the quencher moiety) and a sequence-specific fluorescent signal is generated. With each cycle, additional reporter dye molecules are cleaved from their respective probes, and the fluorescence intensity is monitored at regular intervals by laser optics built into the ABI PRISM™ 7700 Sequence Detection System. In each assay, a series of parallel reactions containing serial dilutions of mRNA from untreated control samples generates a standard curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test samples.

PCR reagents are obtained from PE-Applied Biosystems, Foster City, CA. RT-PCR reactions are carried out by adding 25 μ L PCR cocktail (1x TAQMAN™ buffer A, 5.5 mM $MgCl_2$, 300 μ M each of dATP, dCTP and dGTP, 600 μ M of dUTP, 100 nM each of forward primer, reverse primer, and probe, 20 Units RNase inhibitor, 1.25 Units AMPLITAQ GOLD™, and 12.5 Units MuLV reverse transcriptase) to 96 well plates containing 25 μ L poly(A) mRNA solution. The RT reaction is

-45-

carried out by incubation for 30 minutes at 48°C. Following a 10 minute incubation at 95°C to activate the AMPLITAQ GOLD™, 40 cycles of a two-step PCR protocol are carried out: 95°C for 15 seconds (denaturation) followed by 60°C for 1.5 minutes (annealing/extension). TRAF probes and primers are designed to hybridize to the human TRAF sequence, using published sequence information. For example, GenBank Accession No. U19261, Locus name "HSU 19261" SEQ ID NO: 1; GenBank Accession No. U12597, Locus name "HSU12597" SEQ ID NO. 2; GenBank Accession No U21092, Locus name "HSU21092" SEQ ID NO: 3; GenBank Accession No. X80200, Locus name "HSMLN62" SEQ ID NO. 4; GenBank Accession No. AB000509, Locus name "AB000509" SEQ ID NO. 5; GenBank Accession No. U78798, Locus name "HSU78798" SEQ ID NO. 6.

15 Example 14

Antisense inhibition of TRAF-1 expression- phosphorothioate oligodeoxynucleotides

In accordance with the present invention, a series of oligonucleotides were designed to target different regions of the human TRAF-1 RNA, using published sequences (GenBank accession number U19261, incorporated herein as SEQ ID NO: 1). The oligonucleotides are shown in Table 1. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (GenBank accession no. U19261), to which the oligonucleotide binds. All compounds in Table 1 are oligodeoxynucleotides with phosphorothioate backbones (internucleoside linkages) throughout.

TABLE 1
Nucleotide Sequences of Human TRAF-1 Phosphorothioate
Oligonucleotides

5	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ	TARGET GENE	GENE
		ID	NUCLEOTIDE	TARGET
		NO:	COORDINATES ²	REGION
	26698 TTTAAGTTGCTCCAGGGC	7	0028-0045	5'-UTR
	26699 GCCGGGCGAGGACTGCTG	8	0093-0110	coding
	26700 GCAGACGGTGGGAGGGCA	9	0139-0156	coding
	26701 CTGGGCTCCTTTGGGTCC	10	0159-0176	coding
10	26702 CACAGCAGAGAGCCCTGG	11	0173-0190	coding
	26703 ATTCCTCGGGTTCTCAGA	12	0202-0219	coding
	26704 CCATTCCTCGGGTTCTCA	13	0204-0221	coding
	26705 CCTCGCCATTCTCGGGT	14	0209-0226	coding
	26706 GATCCTCGCCATTCTCG	15	0212-0229	coding
15	26707 AGACGGCTTCCTGGGCTT	16	0270-0287	coding
	26708 TTGAAGGAGCAGCCGACA	17	0351-0368	coding
	26709 GGCCTTCCACTGTTCAT	18	0442-0459	coding
	26710 CCACTTCCACGGCTGCCT	19	0527-0544	coding
	26711 CGCCTGGTGACATTGGTG	20	0894-0911	coding
20	26712 CGCATCATACTCCCCTCT	21	1063-1080	coding
	26713 AGGCGTCAATGGCGTGCT	22	1142-1159	coding
	26714 GGAAGGCGTCAATGGCGT	23	1145-1162	coding
	26715 GGAAGAAGAGTGGGCATC	24	1223-1240	coding
	26716 CGTAGGCGTGCTTGGGTG	25	1259-1276	coding
25	26717 GCCCCGCCCACCCTAAGT	26	1321-1338	stop
	26718 GGAGCCCCGCCACCCTA	27	1324-1341	stop
	26719 CTCAGGAGCCCCGCCAC	28	1328-1345	3'-UTR
	26720 AAGGGCAGGGCATCACAG	29	1380-1397	3'-UTR
	26721 TTTGTGCCCTGAGGTCTT	30	1405-1422	3'-UTR

-47-

	26722	CACCCATCTTTGTGCCCT	31	1413-1430	3'-UTR
	26723	GGCCTCCCAGTGTCGCAT	32	1570-1587	3'-UTR
	26724	CCCGGTCCTGTTTCTGAC	33	1756-1773	3'-UTR
	26725	GCACCCCATCCCTTCCAC	34	1773-1790	3'-UTR
5	26726	TGGAGCCGTCTGGGTTTG	35	1837-1854	3'-UTR
	26727	GTCTTCAAATCCAACCCC	36	1871-1888	3'-UTR
	26728	TTCTGGGCTGGAAGGAAA	37	1896-1913	3'-UTR
	26729	ACTTTCTGGGCTGGAAGG	38	1899-1916	3'-UTR
	26730	AGAGACTTTCTGGGCTGG	39	1903-1920	3'-UTR
10	26731	TTTCCAGAACCCCTGTAG	40	1955-1972	3'-UTR
	26732	ATGTTTCCAGAACCCCTG	41	1958-1975	3'-UTR
	26733	GGGCTGGGTGTGCTCCTG	42	2090-2107	3'-UTR
	26734	TTTATGCCCCTCTTCTTC	43	2204-2221	3'-UTR
	26735	GGAAAGTTTATGCCCCTC	44	2210-2227	3'-UTR
15	26736	TACGGGATTCTGGAAAGC	45	2257-2274	3'-UTR
	26737	AGGTGTTACGGGATTCTG	46	2263-2280	3'-UTR

¹ All cytidines are 5-methyl-cytidines; all linkages are phosphorothioate linkages.

20

² Coordinates from GenBank Accession No. U19261, locus name "HSU19261" SEQ ID NO.1.

Example 15:

25 Antisense inhibition of TRAF-1 expression- phosphorothioate 2'-MOE gapmer oligonucleotides

In accordance with the present invention, a second series of oligonucleotides targeted to human TRAF-1 were synthesized. The oligonucleotide sequences are shown in
 30 Table 2. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (GenBank accession no. U19261), to which the oligonucleotide binds.

-48-

All compounds in Table 2 are chimeric oligonucleotides ("gapmers") 18 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by four-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines.

10

TABLE 2

Nucleotide Sequences of Human TRAF-1 Gapmer
Oligonucleotides

	ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET GENE COORDINATES ²	GENE TARGET REGION
15	26738	TTTAAGTTGCTCCAGGGC	7	0028-0045	5'-UTR
	26739	GCCGGGCGAGGACTGCTG	8	0093-0110	coding
	26740	GCAGACGGTGGGAGGGCA	9	0139-0156	coding
	26741	CTGGGCTCCTTTGGGTCC	10	0159-0176	coding
20	26742	CACAGCAGAGAGCCCTGG	11	0173-0190	coding
	26743	ATTCCTCGGGTTCTCAGA	12	0202-0219	coding
	26744	CCATTCCTCGGGTTCTCA	13	0204-0221	coding
	26745	CCTCGCCATTCCTCGGGT	14	0209-0226	coding
	26746	GATCCTCGCCATTCCTCG	15	0212-0229	coding
25	26747	AGACGGCTTCCTGGGCTT	16	0270-0287	coding
	26748	TTGAAGGAGCAGCCGACA	17	0351-0368	coding
	26749	GGCCTTCCACTGTTTCAT	18	0442-0459	coding
	26750	CCACTTCCACGGCTGCCT	19	0527-0544	coding
	26751	CGCCTGGTGACATTGGTG	20	0894-0911	coding
30	26752	CGCATCATACTCCCCTCT	21	1063-1080	coding
	26753	AGGCGTCAATGGCGTGCT	22	1142-1159	coding

-49-

	26754	GGAAGGCGTCAATGGCGT	23	1145-1162	coding
	26755	GGAAGAAGAGTGGGCATC	24	1223-1240	coding
	26756	CGTAGGCGTGCTTGGGTG	25	1259-1276	coding
	26757	GCCCCGCCCCACCCTAAGT	26	1321-1338	stop
5	26758	GGAGCCCCGCCCCACCCTA	27	1324-1341	stop
	26759	CTCAGGAGCCCCGCCCCAC	28	1328-1345	3'-UTR
	26760	AAGGGCAGGGCATCACAG	29	1380-1397	3'-UTR
	26761	TTTGTGCCCTGAGGTCTT	30	1405-1422	3'-UTR
	26762	CACCCATCTTTGTGCCCT	31	1413-1430	3'-UTR
10	26763	GGCCTCCCAGTGTGCGAT	32	1570-1587	3'-UTR
	26764	CCCGGTCCTGTTTCTGAC	33	1756-1773	3'-UTR
	26765	GCACCCCATCCCTTCCAC	34	1773-1790	3'-UTR
	26766	TGGAGCCGTCTGGGTTTG	35	1837-1854	3'-UTR
	26767	GTCTTCAAATCCAACCCC	36	1871-1888	3'-UTR
15	26768	TTCTGGGCTGGAAGGAAA	37	1896-1913	3'-UTR
	26769	ACTTTCTGGGCTGGAAGG	38	1899-1916	3'-UTR
	26770	AGAGACTTTCTGGGCTGG	39	1903-1920	3'-UTR
	26771	TTTCCAGAACCCCTGTAG	40	1955-1972	3'-UTR
	26772	ATGTTTCCAGAACCCCTG	41	1958-1975	3'-UTR
20	26773	GGGCTGGGTGTGCTCCTG	42	2090-2107	3'-UTR
	26774	TTTATGCCCTCTTCTTC	43	2204-2221	3'-UTR
	26775	GGAAAGTTTATGCCCTC	44	2210-2227	3'-UTR
	26776	TACGGGATTCTGGAAGC	45	2257-2274	3'-UTR
25	26777	AGGTGTTACGGGATTCTG	46	2263-2280	3'-UTR

¹ Emboldened residues are 2'-methoxyethoxy residues (others are 2'-deoxy-). All 2'-methoxyethoxy cytidines and 2'-deoxycytidines are 5-methyl-cytidines; all linkages are phosphorothioate linkages.

-50-

² Coordinates from GenBank Accession No. 19261, locus name "HSU19261" SEQ ID NO. 1.

Example 16:

5 **Antisense inhibition of TRAF-2 expression- phosphorothioate 2'-MOE gapmer oligonucleotides**

10 In accordance with the present invention, a series of oligonucleotides targeted to human TRAF-2 were synthesized. The oligonucleotide sequences are shown in Table 3. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (GenBank accession no. U12597), to which the oligonucleotide binds.

15 All compounds in Table 3 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of eight 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by six-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) in the central "deoxy gap" and phosphodiester (P=O) in the wings. 20 Cytidine residues in the 2'-MOE wings are 5-methylcytidines.

TABLE 3:
Nucleotide Sequences of TRAF-2 Gapmer Oligonucleotides

ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET GENE NUCLEOTIDE COORDINATES ²	GENE TARGET REGION
5				
16827	GoToCoGoCoAsGsCsGsCsGsCsGsGsGoAoAoToToC	47	0001-0020	5'-UTR
16828	CoCoAoAoCoGsGsTsCsGsCsAsGsCsGoCoGoCoCoG	48	0007-0026	5'-UTR
16829	CoAoGoCoCoAsTsGsAsGsCsTsGsToGoAoCoC	49	0042-0061	AUG
16830	AoCoGoCoToAsTsCsTsGsCsAsGsCsCoAoToGoAoG	50	0052-0071	AUG
10				
16831	GoCoCoAoCoAsCsTsGsCsGsCsTsGoGoAoAoGoG	51	0185-0204	coding
16832	CoCoGoGoCoAsGsGsCsTsCsTsCsCsAoCoCoToCoC	52	0348-0367	coding
16833	GoCoAoGoCoGsGsCsTsTsCsGsTsGoGoCoAoGoC	53	0422-0441	coding
16834	CoCoToCoGoTsGsGsTsGsCsGsCsTsToCoAoCoG	54	0576-0595	coding
16835	CoToCoGoAoCsAsCsTsTsGsCsCsAsCoAoAoGoToC	55	0675-0694	coding
15				
16836	CoAoCoToGoCsAsCsTsTsGsTsGsCoToCoCoToG	56	0751-0770	coding
16837	CoCoToCoToGsCsAsGsAsGsCsTsCoToGoAoCoC	57	0848-0867	coding
16838	CoAoGoCoCoGsGsTsGsCsTsGsCsGsGoGoCoToGoC	58	0962-0981	coding
16839	CoCoGoGoToGsCsCsGsTsCsGsCsGsToToCoAoG	59	1240-1259	coding
16840	AoCoGoToCoGsGsGsCsTsGsAsAsGoGoCoGoToC	60	1387-1406	coding

- 52 -

16841	CoToGoToCoAsGsGsTsCsCsAsCsAsAoToGoGoCoC	61	1533-1552	coding
16842	GoCoCoGoGoCsTsGsTsGsCsCsTsGsGoCoToGoCoC	62	1590-1609	3'-UTR
16843	CoToToGoGoCsTsGsCsAsGsGsCsGsAoCoAoCoC	63	1685-1704	3'-UTR
16844	CoGoGoCoCoAsAsTsGsCsCsAsCsAoCoAoGoCoC	64	1789-1808	3'-UTR
16845	AoCoToGoToGsCsTsCsTsGsCsTsAoCoAoToGoG	65	1916-1935	3'-UTR
16846	GoCoToCoToGsGsCsCsAsGsCsAsGsGoAoGoGoCoC	66	1994-2013	3'-UTR
16847	CoCoAoCoAoGsCsCsAsGsCsCsTsGsGoCoCoAoAoG	67	2117-2136	3'-UTR
16848	CoToCoToGoTsCsTsTsCsGsTsGsAsGoCoToGoGoA	68	2221-2240	3'-UTR
26264	CoCoToCoGoTsGsCsTsGsCsGsGsCsToToCoAoCoG	69	mismatch	
26266	CoCoToGoGoTsGsCsTsCsCsGsGsCsToToCoAoCoG	70	mismatch	
27693	CoCoToCoGsTsGsGsTsGsCsGsCsTsToCoAoCoG	54	0576-0595	coding
27694	CsCsTsCsGsTsGsGsTsGsCsGsCsTsTsCsAsCsG	54	0576-0595	coding

¹ Emboldened residues are 2'-methoxyethoxy residues (others are 2'-deoxy). All 2'-methoxyethoxy cytidines are 5-methyl-cytidines, underlined "C" residues are 5-methyl-cytidines; "s" linkages are phosphorothioate linkages, "o" linkages are phosphodiester linkages.

² Coordinates from GenBank Accession No. U12597, locus name "HSU12597" SEQ ID NO.2.

-53-

HMVEC (human dermal microvascular) cells were purchased from Clonetics (San Diego CA) and cultivated in endothelial basal medium (EBM) supplemented with 10% fetal bovine serum (HyClone, Logan UT). Cells were grown in 100 mm petri dishes until 70-80% confluent and then treated with oligonucleotide in the presence of cationic lipid. Briefly, cells were washed with PBS and OPTI-MEM®. OPTI-MEM® containing 10 µg/mL LIPOFECTIN® was added to the cells, followed by addition of oligonucleotide. The cells were incubated for 3-4 hours at 37°C, washed once with EBM/1% FBS, and allowed to recover. For determination of mRNA levels by Northern blot, total RNA was prepared from cells by the guanidinium isothiocyanate procedure or by the Qiagen RNEASY™ method (Qiagen, Valencia, CA). Northern blot analysis was performed by standard methods (for example, Ausubel, et al. *Current Protocols in Molecular Biology*, Vol. 1, John Wiley and Sons, Inc., 1996, pp.4.2.1-4.2.9). The probe was a PCR-labeled 1-kb fragment of TRAF-2 amplified by RT-PCR according to the method of Bednarczuk et al., *Biotechniques*, 1991, 10,478. RNA was quantified and normalized to G3PDH mRNA levels using a Molecular Dynamics (Sunnyvale, CA) PhosphorImager in accordance with manufacturer's instructions.

Results are shown in Table 4. Reduction of TRAF-2 mRNA levels with oligonucleotide 16834 (SEQ ID NO. 54) was determined to be dose-dependent in the range of 1 to 100 nM. The IC₅₀ was approximately 10 nM. A TRAF-6 antisense oligonucleotide did not affect TRAF-2 mRNA expression.

The effect of oligonucleotide 16834 (SEQ ID NO. 54) on TRAF-2 protein levels was also examined. Cells were treated with oligonucleotide and allowed to recover for 48 to 72 hours before being harvested. Protein levels were determined by western blot analysis. Dose-dependent reduction of TRAF-2 protein expression was detectable 48 hours after treatment and maximal reduction of TRAF-2

-54-

protein levels was achieved 72 hours after treatment with 100 nM oligonucleotide.

TABLE 4

5 Activities of TRAF-2 Gapmer Oligonucleotides

	ISIS	SEQ ID	GENE TARGET	% mRNA	% mRNA
	No:	NO:	REGION	EXPRESSION	INHIBITION
10	LIPOFECTIN	---	---	100%	0%
	only				
	16827	47	5'-UTR	43%	57%
	16828	48	5'-UTR	23%	77%
	16829	49	AUG	48%	52%
15	16830	50	AUG	18%	82%
	16831	51	coding	49%	51%
	16832	52	coding	42%	58%
	16833	53	coding	60%	40%
	16834	54	coding	3%	97%
20	16835	55	coding	43%	57%
	16836	56	coding	91%	9%
	16837	57	coding	60%	40%
	16838	58	coding	66%	34%
	16839	59	coding	47%	53%
25	16840	60	coding	45%	55%
	16841	61	coding	8%	92%
	16842	62	3'-UTR	36%	64%
	16843	63	3'-UTR	46%	54%
	16844	64	3'-UTR	82%	18%
30	16845	65	3'-UTR	59%	41%
	16846	66	3'-UTR	13%	87%
	16847	67	3'-UTR	74%	26%
	16848	68	3'-UTR	57%	43%

-55-

ISIS 27693 (SEQ ID NO: 54) was also shown to decrease TRAF-2 mRNA levels in primary human fibroblast-like synoviocytes (obtained from surgical/biopsy specimens). LIPOFECTIN® was included at 3 µg/ml. A dose-response effect was obtained with an IC₅₀ of approximately 25 nM and nearly 90% reduction of TRAF-2 mRNA at an oligonucleotide concentration of 100 nM.

Example 17:

Antisense inhibition of TRAF-3 expression- phosphorothioate oligodeoxynucleotides

In accordance with the present invention, a series of oligonucleotides were designed to target human TRAF-3 RNA using published sequences (GenBank accession number HSU21092, SEQ ID NO: 3). Oligodeoxynucleotides are shown in Table 5. Target sites are indicated as nucleotide numbers on the TRAF-3 mRNA target (SEQ ID NO: 3).

TABLE 5

Nucleotide Sequences of Human TRAF-3 Phosphorothioate Oligonucleotides

ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET GENE COORDINATES ²	GENE TARGET REGION
		ID	NUCLEOTIDE	
20	26778 AGAGCCGACGACCGCCGC	71	0078-0095	5'-UTR
	26779 GGAAGAGCCGACGACCGC	72	0081-0098	5'-UTR
25	26780 CGCGCCAGGAGAGTCCAT	73	0236-0253	coding
	26781 TTAGCGGCGGGTTAGTCT	74	0258-0275	coding
	26782 AGCTTTAGCGGCGGGTTA	75	0262-0279	coding
	26783 CTCGGTCTGCTTCGGGCT	76	0401-0418	coding
	26784 TGCCCACTCGGTCTGC	77	0409-0426	coding
30	26785 CGGTGCCCACACTCGGTC	78	0412-0429	coding
	26786 GAAGCGGTGCCCACACTC	79	0416-0433	coding
	26787 TTACACGCCTTCTCCACG	80	0712-0729	coding

-56-

	26788	GTATTTACACGCCTTCTC	81	0716-0733	coding
	26789	CCGGTATTTACACGCCTT	82	0719-0736	coding
	26790	GAGGGCAGGACACCACCA	83	0816-0833	coding
	26791	TGTGAGGGCAGGACACCA	84	0819-0836	coding
5	26792	CACTTGTTGAGGGCAGGAC	85	0823-0840	coding
	26793	GCTGGTTTGTCCCCCTGAA	86	0939-0956	coding
	26794	ATCTGCTGGTTTGTCCCC	87	0943-0960	coding
	26795	CGCGGTTCTGGAGGGACT	88	1281-1298	coding
	26796	CCCCGCACTCTTGTCCAC	89	1316-1333	coding
10	26797	TTGCCCCGCACTCTTGTC	90	1319-1336	coding
	26798	CCACTTGCCCCGCACTCT	91	1323-1340	coding
	26799	GAGCCACTTGCCCCGCAC	92	1326-1343	coding
	26800	TTCCGAGCCACTTGCCCC	93	1330-1347	coding
	26801	TCCGCCGCTTGTAAGTCGC	94	1485-1502	coding
15	26802	TGCTTCCGCCGCTTGTAAG	95	1489-1506	coding
	26803	TCCTGCTTCCGCCGCTTG	96	1492-1509	coding
	26804	GTCCCCGTTTCAGGTAGAC	97	1589-1606	coding
	26805	TCCCGTCCCCGTTTCAGGT	98	1593-1610	coding
	26806	CCATCCCGTCCCCGTTCA	99	1596-1613	coding
20	26807	TCCCCATCCCGTCCCCGT	100	1599-1616	coding
	26808	CCCTTCCCCATCCCGTCC	101	1603-1620	coding
	26809	TGCGTCCCCCTTCCCCATC	102	1609-1626	coding
	26810	AAGTGCGTCCCCCTTCCCC	103	1612-1629	coding
	26811	CGACAAGTGCGTCCCCCTT	104	1616-1633	coding
25	26812	AAGGAAGCAGGGCATCAT	105	1662-1679	coding
	26813	CTCTCCAGTGGGCTTCTT	106	1781-1798	coding
	26814	TCATCTCTCCAGTGGGCT	107	1785-1802	coding
	26815	GCTAAATCCACCTCCCCA	108	1933-1950	3'-UTR
	26816	TCTGCCGCTTCCTCCGTC	109	2027-2044	3'-UTR
30	26817	CCGCCTTCTGCCGCTTCC	110	2033-2050	3'-UTR

-57-

¹ All cytidines are 5-methyl-cytidines; all linkages are phosphorothioate linkages.

² Coordinates from GenBank Accession No.U21092, locus name "HSU21092" SEQ ID NO.3.

5 Example 18:

Antisense inhibition of TRAF-3 expression- phosphorothioate 2'-MOE gapmer oligonucleotides

In accordance with the present invention, a second series of oligonucleotides targeted to human TRAF-3 were
10 synthesized. The oligonucleotide sequences are shown in Table 6. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (GenBank accession no. U21092), to which the oligonucleotide binds.

All compounds in Table 6 are chimeric oligonucleotides
15 ("gapmers") 18 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by four-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside
20 (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines.

TABLE 6

Nucleotide Sequences of Human TRAF-3 Gapmer 25 Oligonucleotides

ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID	TARGET GENE NUCLEOTIDE COORDINATES ²	GENE TARGET REGION
		NO:		
26818	AGAGCCGACGACCGCCGC	71	0078-0095	5'-UTR
30 26819	GGAAGAGCCGACGACCGC	72	0081-0098	5'-UTR
26820	CGCGCCAGGAGAGTCCAT	73	0236-0253	coding
26821	TTAGCGGCGGGTTAGTCT	74	0258-0275	coding
26822	AGCTTTAGCGGCGGGTTA	75	0262-0279	coding

-58-

	26823	CTCGGTCTGCTTCGGGCT	76	0401-0418	coding
	26824	TGCCCACACTCGGTCTGC	77	0409-0426	coding
	26825	CGGTGCCACACTCGGTC	78	0412-0429	coding
	26826	GAAGCGGTGCCACACTC	79	0416-0433	coding
5	26827	TTACACGCCTTCTCCACG	80	0712-0729	coding
	26828	GTATTTACACGCCTTCTC	81	0716-0733	coding
	26829	CCGGTATTTACACGCCTT	82	0719-0736	coding
	26830	GAGGGCAGGACACCACCA	83	0816-0833	coding
	26831	TGTGAGGGCAGGACACCA	84	0819-0836	coding
10	26832	CACTTGTGAGGGCAGGAC	85	0823-0840	coding
	26833	GCTGGTTTGTCCCTGAA	86	0939-0956	coding
	26834	ATCTGCTGGTTTGTCCCC	87	0943-0960	coding
	26835	CGCGGTTCTGGAGGGACT	88	1281-1298	coding
	26836	CCCCGCACTCTTGTCCAC	89	1316-1333	coding
15	26837	TTGCCCCGCACTCTTGTC	90	1319-1336	coding
	26838	CCACTTGCCCCGCACTCT	91	1323-1340	coding
	26839	GAGCCACTTGCCCCGCAC	92	1326-1343	coding
	26840	TTCCGAGCCACTTGCCCC	93	1330-1347	coding
	26841	TCCGCCGCTTGTAGTCGC	94	1485-1502	coding
20	26842	TGCTTCCGCTCTTGTAG	95	1489-1506	coding
	26843	TCCTGCTTCCGCCGCTTG	96	1492-1509	coding
	26844	GTCCCCGTTTCAGGTAGAC	97	1589-1606	coding
	26845	TCCCGTCCCCGTTTCAGGT	98	1593-1610	coding
	26846	CCATCCCGTCCCCGTTCA	99	1596-1613	coding
25	26847	TCCCCATCCCGTCCCCGT	100	1599-1616	coding
	26848	CCCTTCCCCATCCCGTCC	101	1603-1620	coding
	26849	TGCGTCCCCTTCCCCATC	102	1609-1626	coding
	26850	AAGTGCGTCCCCTTCCCC	103	1612-1629	coding
	26851	CGACAAGTGCGTCCCCTT	104	1616-1633	coding

-59-

	26852	AAGGAAGCAGGGCATCAT	105	1662-1679	coding
	26853	CTCTCCAGTGGGCTTCTT	106	1781-1798	coding
	26854	TCATCTCTCCAGTGGGCT	107	1785-1802	coding
	26855	GCTAAATCCACCTCCCCA	108	1933-1950	3'-UTR
5	26856	TCTGCCGCTTCCTCCGTC	109	2027-2044	3'-UTR
	26857	CCGCCTTCTGCCGCTTCC	110	2033-2050	3'-UTR

¹ Emboldened residues are 2'-methoxyethoxy residues (others are 2'-deoxy-). All 2'-methoxyethoxy cytidines and 2'-
 10 deoxycytidines are 5-methyl-cytidines; all linkages are phosphorothioate linkages.

² Coordinates from GenBank Accession No. U21092, locus name "HSU21092" SEQ ID NO.3.

15

Example 19

Antisense inhibition of TRAF-4 expression- phosphorothioate oligodeoxynucleotides

In accordance with the present invention, a series of
 20 oligonucleotides were designed to target different regions of the human TRAF-4 RNA, using published sequences (GenBank accession number X80200, incorporated herein as SEQ ID NO: 4). The oligonucleotides are shown in Table 7. Target sites are indicated by nucleotide numbers, as given in the
 25 sequence source reference (GenBank accession no. X80200), to which the oligonucleotide binds. All compounds in Table 7 are oligodeoxynucleotides with phosphorothioate backbones (internucleoside linkages) throughout. The compounds are analyzed for effect on TRAF mRNA levels by quantitative
 30 real-time PCR as described in other examples herein.

-60-

TABLE 7

Nucleotide Sequences of Human TRAF-4 Phosphorothioate
Oligonucleotides

			SEQ	TARGET GENE	GENE
	ISIS	NUCLEOTIDE SEQUENCE ¹	ID	NUCLEOTIDE	TARGET
	NO.	(5' -> 3')	NO:	COORDINATES ²	REGION
5	26860	GCATGGCGGGCGAGCGGC	111	0072-0089	AUG
	26861	CCGTCGCTTGGGCTTCTC	112	0113-0130	coding
	26862	GGGCACTTGAAGACTCCT	113	0232-0249	coding
10	26863	CTCAGGGCACTTGAAGAC	114	0236-0253	coding
	26864	TGGTCCTCAGGGCACTTG	115	0241-0258	coding
	26865	AAGCTGGTCCTCAGGGCA	116	0245-0262	coding
	26866	GCGGCAGCCCTCCTCACT	117	0341-0358	coding
	26867	CTCCAGCGGCAGCCCTCC	118	0346-0363	coding
15	26868	TAGGGCAGGGAATGACAT	119	0411-0428	coding
	26869	CGATTAGGGCAGGGAATG	120	0415-0432	coding
	26870	GGGCAGCGATTAGGGCAG	121	0421-0438	coding
	26871	GCCTCCCCACTGAAGTCA	122	0523-0540	coding
	26872	ATGCGGGCACCACACTTA	123	0592-0609	coding
20	26873	GGGCAGGCAACAGGCAGC	124	0733-0750	coding
	26874	CCACAGTGCCACACCAC	125	0759-0776	coding
	26875	CGAGCCACAGTGCCACACA	126	0763-0780	coding
	26876	TCCTCCCGAGCCACAGTG	127	0769-0786	coding
	26877	CAGGTCCTCCCGAGCCAC	128	0773-0790	coding
25	26878	GGCAGAGCACCAGGGCGG	129	0819-0836	coding
	26879	CTTTGAATGGGCAGAGCA	130	0828-0845	coding
	26880	GGAGTCTTTGAATGGGCA	131	0833-0850	coding
	26881	ATGCCGTGCCATTGCCAG	132	0875-0892	coding
	26882	CTCACCAGGGCACACATC	133	0925-0942	coding
30	26883	CAGCTCCTGCCGTTGCCG	134	0944-0961	coding
	26884	ATGAGCACGCCATCACTG	135	1000-1017	coding

-61-

	26885	TGTAGCCGCCGTCCATAG	136	1033-1050	coding
	26886	GCCTCCTGTAGCCGCCGT	137	1039-1056	coding
	26887	TAGAAGGCTGGGCTGAAG	138	1081-1098	coding
	26888	GTGTGTAGAAGGCTGGGC	139	1086-1103	coding
5	26889	GTGTGCCCTCACCCTGTC	140	1152-1169	coding
	26890	GACACGGCGGGCAAAGGG	141	1226-1243	coding
	26891	GAAGGTGACACGGCGGGC	142	1232-1249	coding
	26892	GCCCAGGGTCGCTCTGAT	143	1260-1277	coding
	26893	CTTCCAGTTTGGGTCGGG	144	1313-1330	coding
10	26894	GATAACCAAAGCCCAGAG	145	1377-1394	coding
	26895	CATCGTCCTTTCCCCTCG	146	1513-1530	3'-UTR
	26896	GGCCAGGGCTGAAGCACC	147	1660-1677	3'-UTR
	26897	TTGTTTCCAGCCCTTCAT	148	1703-1720	3'-UTR
	26898	CATGTCTGCCCTACCCAA	149	1746-1763	3'-UTR
15	26899	GCTCCCCTGCTGTGCCCT	150	1948-1965	3'-UTR

¹ All cytidines are 5-methyl-cytidines; all linkages are phosphorothioate linkages.

20 ² Coordinates from GenBank Accession No. X80200, locus name "HSMLN62" SEQ ID NO. 4.

Example 20:

Antisense inhibition of TRAF-4 expression- phosphorothioate

25 2'-MOE gapmer oligonucleotides

In accordance with the present invention, a second series of oligonucleotides targeted to human TRAF-4 were synthesized. The oligonucleotide sequences are shown in Table 8. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (GenBank
30 accession no. X80200), to which the oligonucleotide binds.

All compounds in Table 8 are chimeric oligonucleotides ("gapmers") 18 nucleotides in length, composed of a central

-62-

"gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by four-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines.

Data are obtained by real-time quantitative PCR as described in other examples herein.

10

TABLE 8

Nucleotide Sequences of Human TRAF-4 Gapmer
Oligonucleotides

	ISIS	NUCLEOTIDE SEQUENCE ¹	SEQ ID	TARGET GENE	GENE
	NO.	(5' -> 3')	NO:	COORDINATES ²	TARGET REGION
15	26900	GCATGGCGGGCGAGCGGC	111	0072-0089	AUG
	26901	CCGTCGCTTGGGCTTCTC	112	0113-0130	coding
	26902	GGGCACTTGAAGACTCCT	113	0232-0249	coding
	26903	CTCAGGGCACTTGAAGAC	114	0236-0253	coding
20	26904	TGGTCCTCAGGGCACTTG	115	0241-0258	coding
	26905	AAGCTGGTCCTCAGGGCA	116	0245-0262	coding
	26906	GCGGCAGCCCTCCTCACT	117	0341-0358	coding
	26907	CTCCAGCGGCAGCCCTCC	118	0346-0363	coding
	26908	TAGGGCAGGGAATGACAT	119	0411-0428	coding
25	26909	CGATTAGGGCAGGGAATG	120	0415-0432	coding
	26910	GGGCAGCGATTAGGGCAG	121	0421-0438	coding
	26911	GCCTCCCCACTGAAGTCA	122	0523-0540	coding
	26912	ATGCGGGCACCACACTTA	123	0592-0609	coding
	26913	GGGCAGGCAACAGGCAGC	124	0733-0750	coding
30	26914	CCACAGTGCCACACCAC	125	0759-0776	coding
	26915	CGAGCCACAGTGCCACACA	126	0763-0780	coding

-63-

	26916	TCCTCCCGAGCCACAGTG	127	0769-0786	coding
	26917	CAGGTCCTCCCGAGCCAC	128	0773-0790	coding
	26918	GGCAGAGCACCAGGGCGG	129	0819-0836	coding
	26919	CTTTGAATGGGCAGAGCA	130	0828-0845	coding
5	26920	GGAGTCTTTGAATGGGCA	131	0833-0850	coding
	26921	ATGCCGTGCCATTGCCAG	132	0875-0892	coding
	26922	CTCACCAGGGCACACATC	133	0925-0942	coding
	26923	CAGCTCCTGCCGTTGCCG	134	0944-0961	coding
	26924	ATGAGCACGCCATCACTG	135	1000-1017	coding
10	26925	TGTAGCCGCCGTCCATAG	136	1033-1050	coding
	26926	GCCTCCTGTAGCCGCCGT	137	1039-1056	coding
	26927	TAGAAGGCTGGGCTGAAG	138	1081-1098	coding
	26928	GTGTGTAGAAGGCTGGGC	139	1086-1103	coding
	26929	GTGTGCCCTCACCCTGTC	140	1152-1169	coding
15	26930	GACACGGCGGGCAAAGGG	141	1226-1243	coding
	26931	GAAGGTGACACGGCGGGC	142	1232-1249	coding
	26932	GCCCAGGGTCGCTCTGAT	143	1260-1277	coding
	26933	CTTCCAGTTTGGGTGCGG	144	1313-1330	coding
	26934	GATAACCAAAGCCCAGAG	145	1377-1394	coding
20	26935	CATCGTCCATTCCCTCG	146	1513-1530	3'-UTR
	26936	GGCCAGGGCTGAAGCACC	147	1660-1677	3'-UTR
	26937	TTGTTTCCAGCCCTTCAT	148	1703-1720	3'-UTR
	26938	CATGTCTGCCCTACCCAA	149	1746-1763	3'-UTR
25	26939	GCTCCCCTGCTGTGCCCT	150	1948-1965	3'-UTR

¹ Emboldened residues are 2'-methoxyethoxy residues (others are 2'-deoxy-). All 2'-methoxyethoxy cytidines and 2'-deoxycytidines are 5-methyl-cytidines; all linkages are phosphorothioate linkages.

-64-

² Coordinates from GenBank Accession No. X80200, locus name "HSMLN62" SEQ ID NO. 4.

Example 21

5 **Antisense inhibition of TRAF-5 expression- phosphorothioate oligodeoxynucleotides**

 In accordance with the present invention, a series of oligonucleotides were designed to target different regions of the human TRAF-5 RNA, using published sequences (GenBank
10 accession number AB000509, incorporated herein as SEQ ID NO: 5). The oligonucleotides are shown in Table 9. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (GenBank accession no. AB000509), to which the oligonucleotide binds. All compounds in Table
15 9 are oligodeoxynucleotides with phosphorothioate backbones (internucleoside linkages) throughout.

TABLE 9
Nucleotide Sequences of Human TRAF-5 Phosphorothioate Oligonucleotides

ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET GENE NUCLEOTIDE COORDINATES ²	GENE TARGET REGION
5	26940 TGAATAAGCCATTGTGGG	151	0049-0066	AUG
	26941 CTTTATGCTCTTCTGAAT	152	0062-0079	coding
	26942 GGATGAAACCACAGGGCA	153	0083-0100	coding
	26943 TCAAAGTCCAAGGAAATG	154	0120-0137	coding
	26944 TGAAGCACCGAGTGGCAG	155	0195-0212	coding
10	26945 GGGCAGATTGGCACTGTG	156	0282-0299	coding
	26946 CTCCTGAGATTGTGATGAC	157	0313-0330	coding
	26947 CTTCCGCTAGGACTGGCT	158	0491-0508	coding
	26948 GATTCTGTAGATTGATGA	159	0584-0601	coding
	26949 TTCATCTACCTCAGTTTT	160	0667-0684	coding
15	26950 TCCGTTACAGCACAGCCA	161	0735-0752	coding
	26951 GCATGTGCTCCCGTAAGG	162	0788-0805	coding
	26952 CTTTTCAAGTTTCTTTAT	163	0907-0924	coding
	26953 CTTCCATCAAAGGTCTCA	164	1079-1096	coding

26954	TCTAAAACGGCTAATCTT	165	1146-1163	coding
26955	TCATCTTGTAATCTGTCA	166	1283-1300	coding
26956	GGA CTGGCTGAAGATGGA	167	1333-1350	coding
26957	CCCTCCCTGACCCATCCC	168	1403-1420	coding
26958	GAATGAGCCACAAAAGCGG	169	1620-1637	coding
26959	CAAGAACAGAGTGTCA TC	170	1672-1689	coding
26960	GTCTAAATCCAGGTCAAT	171	1799-1816	3'-UTR
26961	AAACTTACCATCTTTCAA	172	1964-1981	3'-UTR
26962	CTCTGTGTCTCCATAAC	173	2053-2070	3'-UTR
26963	CTTAACTGGAACAGCCTA	174	2167-2184	3'-UTR
26964	GCAGGAAGAATGAAAAATG	175	2352-2369	3'-UTR
26965	TATTGGTTGAATCTTAT	176	2501-2518	3'-UTR
26966	AAATCTATCCATCTCTCA	177	2611-2628	3'-UTR
26967	AAATTGTAAAGGTTTCT	178	2683-2700	3'-UTR
26968	ACAATGAAACTCTGTCTC	179	2779-2796	3'-UTR
26969	GCAAACTCCGTCTCTAC	180	2940-2957	3'-UTR
26970	CAATAGTTGTCAGAGGCT	181	3055-3072	3'-UTR
26971	AAGGACTCATCTCAGTTT	182	3209-3226	3'-UTR
26972	TAACAACGCAGAAGGGCT	183	3280-3297	3'-UTR

26973	AGTAGGGAAGTGGCATAA	184	3295-3312	3'-UTR
26974	CATCACCAGGTAAGCAGC	185	3377-3394	3'-UTR
26975	TCCTGTTGTGAACCTATT	186	3553-3570	3'-UTR
26976	GGACTTGTGGGCTAAAAGA	187	3656-3673	3'-UTR
26977	GCTCAGGAAGACAGAGTG	188	3724-3741	3'-UTR
26978	TGAACTCCTAAGCAAACC	189	3873-3890	3'-UTR
26979	GATGATGAAGGAACTCTG	190	3889-3906	3'-UTR

¹ All cytidines are 5-methyl-cytidines; all linkages are phosphorothioate linkages.

² Coordinates from GenBank Accession No. AB000509, locus name "AB000509" SEQ ID NO. 5.

-68-

Example 22:**Antisense inhibition of TRAF-5 expression- phosphorothioate 2'-MOE gapmer oligonucleotides**

In accordance with the present invention, a second
5 series of oligonucleotides targeted to human TRAF-5 were
synthesized. The oligonucleotide sequences are shown in
Table 10. Target sites are indicated by nucleotide
numbers, as given in the sequence source reference (GenBank
accession no. AB000509), to which the oligonucleotide
10 binds.

All compounds in Table 10 are chimeric
oligonucleotides ("gapmers") 18 nucleotides in length,
composed of a central "gap" region consisting of ten 2'-
deoxynucleotides, which is flanked on both sides (5' and 3'
15 directions) by four-nucleotide "wings". The wings are
composed of 2'-methoxyethyl (2'-MOE)nucleotides. The
internucleoside (backbone) linkages are phosphorothioate
(P=S) throughout the oligonucleotide. All cytidine
residues are 5-methylcytidines.

TABLE 10
Nucleotide Sequences of Human TRAF-5 Gapmer Oligonucleotides

5	ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID	TARGET GENE NUCLEOTIDE COORDINATES ²	GENE TARGET REGION
10	26980	TGAATAAGCCATTGTGGG	151	0049-0066	AUG
	26981	CTTTATGCTCTTCTGAAT	152	0062-0079	coding
	26982	GGATGAACACACAGGGCA	153	0083-0100	coding
	26983	TCAAAGTCCAAGGAATG	154	0120-0137	coding
	26984	TGAAGCACCGAGTGGCAG	155	0195-0212	coding
	26985	GGGCAGATTGGCACTGTG	156	0282-0299	coding
	26986	CTCCTGAGATTTGATGAC	157	0313-0330	coding
	26987	CTTCCGTAGGACTGGCT	158	0491-0508	coding
	26988	GATTCTGTAGATTGATGA	159	0584-0601	coding
	26989	TTCATCTACCTCAGTTT	160	0667-0684	coding
15	26990	TCCGTTACAGCACAGCCA	161	0735-0752	coding
	26991	GCATGTGCTCCCGTAAGG	162	0788-0805	coding
	26992	CTTTTCAAGTTTCTTTAT	163	0907-0924	coding

26993	CTTCCATCAAAGGTCTCA	164	1079-1096	coding
26994	TCTAAACCGGCTAATCTT	165	1146-1163	coding
26995	TCATCTTGTAATCTGTCA	166	1283-1300	coding
26996	GGACTGGCTGAAGATGGA	167	1333-1350	coding
26997	CCCTCCCTGACCCATCCC	168	1403-1420	coding
26998	GAATGAGCCACAAAGCGG	169	1620-1637	coding
26999	CAAGAACAGAGTGTCAATC	170	1672-1689	coding
27000	GTCTAAATCCAGGTCAAT	171	1799-1816	3'-UTR
27001	AAACTTACCATCTTTCAA	172	1964-1981	3'-UTR
27002	CTCTGTGTCTCTCCATAAC	173	2053-2070	3'-UTR
27003	CTTAACTGGAACAGCCTA	174	2167-2184	3'-UTR
27004	GCAGGAAGAATGAAAATG	175	2352-2369	3'-UTR
27005	TATTTGGTTGAATCTTAT	176	2501-2518	3'-UTR
27006	AAATTCATCCATCCTCA	177	2611-2628	3'-UTR
27007	AAATTGTAAAGGTTTTCT	178	2683-2700	3'-UTR
27008	ACAATGAAACTCTGTCTC	179	2779-2796	3'-UTR
27009	GCAAAACTCCGCTCTAC	180	2940-2957	3'-UTR
27010	CAATAGTTGTCAGAGGCT	181	3055-3072	3'-UTR

- 71 -

27011	AAGGACTCATCTCAGTTT	182	3209-3226	3'-UTR
27012	TAACAACGCAGAAGGGCT	183	3280-3297	3'-UTR
27013	AGTAGGGAAGTGGCATAA	184	3295-3312	3'-UTR
27014	CATCACCAGGT/ AGCAGC	185	3377-3394	3'-UTR
27015	TCCTGTTGTGAACCTATT	186	3553-3570	3'-UTR
27016	GGACTTGTGGGCTAAAGA	187	3656-3673	3'-UTR
27017	GCTCAGGAAGACAGAGTG	188	3724-3741	3'-UTR
27018	TGAACTCCTAAGCAAACC	189	3873-3890	3'-UTR
27019	GATGATGAAGGAACTCTG	190	3889-3906	3'-UTR

10

¹ Emboldened residues are 2'-methoxyethoxy residues (others are 2'-deoxy-). All 2'-methoxyethoxy cytidines and 2'-deoxycytidines are 5-methyl-cytidines; all linkages are phosphorothioate linkages.

² Coordinates from GenBank Accession No. AB000509, locus name "AB000509" SEQ ID NO. 5.

-72-

Example 23:**Antisense inhibition of TRAF-6 expression- phosphorothioate 2'-MOE gapmer oligonucleotides**

In accordance with the present invention, a series of
5 oligonucleotides targeted to human TRAF-6 were synthesized.
The oligonucleotide sequences are shown in Table 11.
Target sites are indicated by nucleotide numbers, as given
in the sequence source reference (GenBank accession no.
U78798), to which the oligonucleotide binds.

10 All compounds in Table 11 are chimeric
oligonucleotides ("gapmers") 20 nucleotides in length,
composed of a central "gap" region consisting of eight 2'-
deoxynucleotides, which is flanked on both sides (5' and 3'
directions) by six-nucleotide "wings". The wings are
15 composed of 2'-methoxyethyl (2'-MOE)nucleotides. The
internucleoside (backbone) linkages are phosphorothioate
(P=S) in the "deoxy gap" and phosphodiester (P=O) in the
wings. Cytidine residues in the 2'-MOE wings are 5-
methylcytidines.

20

TABLE 11:
Nucleotide Sequences of TRAF-6 Gapmer Oligonucleotides

ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET GENE	
			NUCLEOTIDE COORDINATES ²	TARGET REGION
5	15779 AOGGOCOCOAsAGSCsCsAGSCsCOTOGOCOGG	191	0001-0020	5'-UTR
	15880 CCGOCOCOAoCsCstTsCsGSCstTsGSCoCoGoCoC	192	0024-0043	5'-UTR
	15881 GAOGoAOCoGsAGSGsCstTsGSCstTsTOGoGoAOCoG	193	0071-0090	5'-UTR
	15882 GOGAOCoAOCSAGSAsCsAsCstTsGSCOGOCOGCoC	194	0091-0110	5'-UTR
	15883 COCoAOAGOGGSCSGsCstTsGSGstTsAGoAGOGGoAOC	195	0111-0130	5'-UTR
10	15884 TOTOGoCoToCsGstTsCsTsAGstTsGOCOGOCOGG	196	0185-0204	5'-UTR
	15885 CoAoToAOGoTsAsAsCstTsGstTsAGstTsAoToCoAoC	197	0205-0224	AUG
	15886 AOGCoAOGoAsCstTsCsAsTsAGstTsAoAOCoToToG	198	0213-0232	AUG
	15887 AOCoAOGoToTsTsAGsCsAGsAsCstToCoAoToAoG	199	0220-0239	AUG
	15888 AOCoAOGoCoGsCstTsAsCsAGsGstTsGOCOGOC	200	0291-0310	coding
15	15889 AOToTOGAOTsTsTsAsTsGstTsAGstTsGOCOGOGOC	201	0495-0514	coding
	15890 GOTOGAOCOCsTsGSCsAsTsCsCsCstToAoToToG	202	0511-0530	coding
	15891 GOTOCoToCoAGstTsCsCsAsTsCsToToGoToGoC	203	0641-0660	coding
	15892 AOGAOGoCoAsAsCstTsCsAsCsAoToGoToGoC	204	0678-0697	coding

15893	TtTtTtTtGgGgAaAaGgGgAaCsGsCoToGoGoCoA	205	0714-0733	coding
15894	AaAaCoToGoCoCsAaTtTtGgAaTtGgCoAoGoCoAoC	206	0796-0815	coding
15895	AaToToCoAoCsAaGgAaTtGgAaCsAaTtToToGoCoC	207	0851-0870	coding
15896	CoGoToGoCoCsAaAaGgTtGgAaTtTsCoCoToCoToG	208	0981-1000	coding
15897	GoGoToGoToTsCsTsCsTsTsGgTsAaGoGoToGoGoC	209	1000-1019	coding
15898	GoGoCoCoAoAaCsAaTtTsCsTsCsAaToGoToGoToG	210	1024-1043	coding
15899	CoGoCoToCoAaAaCsAaTtTsAaTtGgAaCoAoGoCoC	211	1046-1065	coding
15900	AaGoGoCoGoAaCsCsCsTsCsTsAaCoToGoGoToG	212	1119-1138	coding
15901	CoCoAoToToTsTsAaGgCsAaGgTsCsAaGoCoToCoC	213	1163-1182	coding
15902	CoGoAoAoToGgGgTsTsCsGgTsTsTsGoAoGoCoToC	214	1206-1225	coding
15903	CoCoAoToToGgCsAaCsTsGsCsTsGsToGoCoToToC	215	1254-1273	coding
15904	GoCoAoGoToCsGgTsAaAaCsTsGsAaAoGoGoToG	216	1401-1420	coding
15905	GoCoCoToToAaCsAaGgGgTsGsCsTsToCoAoGoAoC	217	1532-1551	coding
15906	AaGoCoAoAoGgCsAaGgCsTsCsTsGsGoToToToGoG	218	1576-1595	coding
15907	GoGoCoToAoCsCsAaTtGgTsCsAaAoGoCoGoG	219	1724-1743	coding
15908	ToToGoToToToTsTsTsGsAaGgCsAaGoToGoAoGoG	220	1796-1815	3'-UTR
15909	GoGoCoAoCoTsGgTsTsTsCsTsCsCoAoGoGoToA	221	1817-1836	3'-UTR
15910	AaCoAoToAoTsTsCsCsGsTsGsGoCoToToGoT	222	1871-1890	3'-UTR

15911	GoGoAoAoCoG TsTsGsTsGsAsTsTsCoCoCoAoGoG	223	1967-1986	3'-UTR
15912	ToGoCoToGoCsAsAsCsAsTsGsCsCsAoCoAoGoGoC	224	2017-2036	3'-UTR
15913	AoToAoCoAoCsAsGsAsGsCsAsAsAoAoGoCoCoC	225	2078-2097	3'-UTR
15914	AoAoAoGoAsCsTsGsAsCsTsTsToToAoAoGoG	226	scrambled	
			control	
5	23247	AoCoToToAoAsTsTsAsCsCsAsTsGsAoCoToAoGoT	227	15910 mismatch
	23248	CoCoAoCoGoAsGsGsAsGsCsAsCsAoToCoAoAoG	228	16834 mismatch
	27691	AsCsAsTsAsTsTsTsCsCsGsTsGsGsCsTsTsGst	222	1871-1890
	27692	AoCoAoToAsTsTsTsCsCsGsTsGsGsCsToToToGoT	222	1871-1890
				3'-UTR
				3'-UTR

- 75 -

¹ Emboldened residues are 2'-methoxyethoxy residues (others are 2'-deoxy). All 2'-methoxyethoxy cytidines are 5-methyl-cytidines, underlined "C" are 5-methyl-cytidine; "s" linkages are phosphorothioate linkages, "o" linkages are phosphodiester linkages.

² Coordinates from GenBank Accession No. U78798, locus name "HSU78798" SEQ ID NO. 6.

-76-

HMVEC cells were grown in 100 mm petri dishes until 70-80% confluent and then treated with oligonucleotide in the presence of cationic lipid. Briefly, cells were washed with PBS and OPTI-MEM®. OPTI-MEM® containing 10 µg/mL LIPOFECTIN® (Life Technologies, Rockville MD) was added to the cells, followed by addition of oligonucleotide. The cells were incubated for 3-4 hours at 37°C, washed once with EBM/1% FBS, and allowed to recover. For determination of mRNA levels by Northern blot, total RNA was prepared from cells by the guanidinium isothiocyanate procedure or by the Qiagen RNEASY™ method (Qiagen, Valencia, CA). Northern blot analysis was performed by standard methods (for example, Ausubel, et al. *Current Protocols in Molecular Biology*, Vol. 1, John Wiley and Sons, Inc., 1996, pp.4.2.1-4.2.9). The probe was a PCR-labeled 1-kb fragment of TRAF-6 amplified by RT-PCR according to the method of Bednarczuk et al., 1991, *Biotechniques* 10,478. RNA was quantified and normalized to G3PDH mRNA levels using a Molecular Dynamics (Sunnyvale, CA) PhosphorImager in accordance with manufacturer's instructions.

Results are shown in Table 12. Reduction of TRAF-6 mRNA levels with oligonucleotide 15910 (SEQ ID NO. 224) was determined to be dose-dependent in the range of 1 to 100 nM. The IC₅₀ was approximately 2.5 nM. A TRAF-2 antisense oligonucleotide did not affect TRAF-6 mRNA expression.

TABLE 12

Activities of TRAF-6 Gapmer Oligonucleotides

	ISIS	SEQ	GENE	% mRNA	% mRNA
	No:	ID	TARGET	EXPRESSION	INHIBITION
		NO:	REGION		
30	LIPOFECTIN®	---	---	100%	0%
	only				
	15779	191	5'-UTR	62%	38%

-77-

	15880	192	5'-UTR	73%	27%
	15881	193	5'-UTR	28%	72%
	15882	194	5'-UTR	96%	4%
	15883	195	5'-UTR	57%	43%
5	15884	196	5'-UTR	73%	27%
	15885	197	AUG	61%	39%
	15886	198	AUG	37%	63%
	15887	199	AUG	23%	77%
	15888	200	coding	31%	69%
10	15889	201	coding	42%	58%
	15890	202	coding	49%	51%
	15891	203	coding	50%	50%
	15892	204	coding	32%	68%
	15893	205	coding	18%	82%
15	15894	206	coding	43%	57%
	15895	207	coding	41%	59%
	15896	208	coding	20%	80%
	15897	209	coding	60%	40%
	15898	210	coding	23%	77%
20	15899	211	coding	66%	34%
	15900	212	coding	54%	46%
	15901	213	coding	60%	40%
	15902	214	coding	76%	24%
	15903	215	coding	58%	42%
25	15904	216	coding	77%	23%
	15905	217	coding	108%	---
	15906	218	coding	90%	10%
	15907	219	coding	62%	38%
	15908	220	3'-UTR	82%	18%
30	15909	221	3'-UTR	28%	72%

-78-

	15910	222	3'-UTR	13%	87%
	15911	223	3'-UTR	103%	---
	15912	224	3'-UTR	20%	80%
	15913	225	3'-UTR	97%	3%
5	15914	226	scrambled control	70%	30%

Example 24

Effect of inhibiting TRAF Gene Expression on the Induction of E-selectin

10 The effect of TRAF antisense oligonucleotides on the induction of E-selectin by $\text{TNF}\alpha$ or $\text{IL-1}\beta$ was examined. HMVEC cells were treated with either ISIS 16834 or ISIS 15910 under dose-response conditions followed by stimulation of E-selectin expression by $\text{TNF}\alpha$ or $\text{IL-1}\beta$ for 5

15 hours. The cell surface expression of E-selectin was determined by flow cytometry analysis. Dose-dependent inhibition of E-selectin cell surface induction by $\text{TNF}\alpha$ was observed in cells treated with the TRAF-2 antisense oligonucleotide ISIS 16834, as expected. Surprisingly, the

20 TRAF-6 antisense compound, ISIS 15910, was able to inhibit $\text{TNF}\alpha$ mediated E-selectin surface expression as well, especially at higher dose. At low doses (20-50nM), ISIS 16834 was a more effective inhibitor of $\text{TNF}\alpha$ -mediated E-selectin induction than ISIS 15910. Maximal inhibition of

25 E-selectin induction for both antisense compounds was approximately 70% at 100 nM. Control oligonucleotides exhibited little to no effect on E-selectin induction. When $\text{IL-1}\beta$ was used as the stimulator, however, ISIS 15910 appeared to be a more specific and potent inhibitor of E-

30 selectin induction than ISIS 16834, especially at relatively low doses.

-79-

Example 25**Effect of TRAF Antisense Oligonucleotide on I κ B α Phosphorylation and Degradation**

Multiple transcription factors are activated by cytokines to facilitate the induction of E-selectin. The most important and best studied transcription factors involved in the regulation of E-selectin activation include NF- κ B, c-Jun and ATF-2. To clarify the roles of TRAF proteins in the activation of NF- κ B by cytokines, I κ B α phosphorylation and degradation assays were performed with antisense oligonucleotide treated cells. Cells were treated with either ISIS 16834 or ISIS 15910 and allowed to recover for 48-72 hours. Tumor necrosis factor- α (TNF- α) or interleukin-1- β (IL-1 β) was added for 5 to 30 minutes before cells were harvested. Western blot analysis with antibody specific for phospho-I κ B α was performed to study the phosphorylation of I κ B α . The blots were then stripped and reblotted with antibody against I κ B α to study the degradation of I κ B α . I κ B α was heavily phosphorylated 5 minutes after addition of either cytokine. By 30 minutes, I κ B α was reduced, probably as a result of I κ B α degradation. In TNF α -stimulated cells, the majority of the I κ B α had been degraded after 5 minutes of stimulation. By 30 minutes, I κ B α was almost completely gone. In contrast, the degradation of I κ B α in IL-1 β stimulated cells was slower with the majority of I κ B α degraded by 30 minutes. Neither ISIS 16834 nor ISIS 15910 affected I κ B α phosphorylation and degradation induced by TNF α . ISIS 15910 has little effect on IL-1 β mediated I κ B α phosphorylation and degradation either. Hyperphosphorylation of I κ B α was observed in ISIS 16834 treated, IL-1 β induced cells. In summary, the antisense oligonucleotides do not inhibit the phosphorylation and degradation of I κ B α .

-80-

Example 26**Effect of TRAF Antisense Oligonucleotides on JNK Activities**

MAP kinases play central roles in the activation of specific transcription factors crucial to the induction of cell adhesion molecules. To examine the effect of TRAF antisense oligonucleotides on JNK activities, *in vitro* kinase assays were performed on extracts derived from cells treated with TRAF antisense oligonucleotides. Cells were treated with TRAF-2 or TRAF-6 antisense compounds, (ISIS 16834 or ISIS 15910, respectively) allowed to recover for 48-72 hours, at which time TNF was added for 15 minutes prior to the cell lysis and the initiation of the kinase assays. Specific c-Jun conjugated agarose beads were used to precipitate JNK. ATP was added to the immunoprecipitated kinase complexes and the reaction mixtures were analyzed on SDS-PAGE. Western blotting with antibodies specific for phosphorylated c-Jun was carried out to determine relative kinase activity. JNK was activated by TNF α after a 15 minute incubation, as indicated by the heavy phosphorylation of c-Jun. ISIS 16834 reduced JNK activity in TNF α -treated cells but not in IL-1 β treated cells. Some hyperphosphorylation of c-Jun induced by IL-1 β in ISIS 16834 treated cells was observed. ISIS 15910 reduced the c-Jun phosphorylation mediated by both IL-1 β and TNF α . Some inhibitory effect of ISIS 15910 on JNK activity was also observed in TNF α -induced cells. This result is consistent with the inhibitory effects of TRAF antisense oligonucleotides on the surface expression of E-selectin.

Example 27**Inhibition of cell proliferation by antisense oligonucleotide targeted to TRAF-2**

HeLa cells were treated with ISIS 16834 (200 nM) and counted 48 hours later. Cells were trypsinized, stained

-81-

with trypan blue and counted (floating cells included). Cell number in ISIS 16834-treated cultures was reduced by 61% compared to untreated control cultures.

Example 28

- 5 Induction of apoptosis by antisense oligonucleotide targeted to TRAF-2

HeLa cells were treated with ISIS 16834 (200 nM) and the number of dead cells was measured by trypan blue exclusion 48 hours later. In cultures untreated with
10 oligonucleotide, only 5% of cells were dead. In cultures treated with ISIS 16834, 44% of cells were dead.

Example 29

Dose response of apoptosis in response to ISIS 16834 targeted to TRAF-2

- 15 HeLa cells were treated with ISIS 16834 at various doses and the number of sub-G1 apoptotic cells was counted using propidium iodide and FACS 48 hours after treatment with ISIS 16834 at 100, 200 and 300 nM. Culture supernatant and floating cells were transferred to FACS tubes. Cells
20 were washed with PBS, trypsinized and washed in PBS, then fixed in ice-cold 70% ethanol for 12-15 hours in the freezer. Cells were centrifuged and resuspended in propidium iodide (PI) mix (50 µg/ml PI, 5 µg/ml RNase cocktail, Cat. # 2286, Ambion, Austin TX) in the dark for
25 1 hour at room temperature before FACS analysis. Percent apoptotic cells after treatment at 100, 200 and 300 nM doses was approximately 29%, 46% and 58%, respectively, compared to 5% for untreated control cells.

Example 30

- 30 Time course of apoptosis in response to ISIS 16834 targeted to TRAF-2

The number of sub-G1 apoptotic HeLa cells was counted 1, 2 and 3 days after treatment with ISIS 16834 at 200 nM using methods described in the previous example. Percent

-82-

apoptotic cells after treatment at these time points was approximately 10%, 46% and 53%, respectively, compared to 5% after 3 days for untreated control cells.

Example 31**5 Antisense inhibition of TRAF-3 expression**

The antisense oligonucleotides shown in Tables 5 and 6 were screened in T-24 cells for ability to inhibit human TRAF-3 expression. Results are shown in Tables 13 and 14, respectively.

-83-

Table 13

Inhibition of TRAF-3 mRNA levels by phosphorothioate
oligodeoxynucleotides

	ISIS#	TARGET REGION	SITE ¹	SEQUENCE 5'--> 3'	%Inhib	SEQ ID NO.
5	26778	5'UTR	78	AGAGCCGACGACCGCCGC	0	71
	26779	5'UTR	81	GGAAGAGCCGACGACCGC	0	72
	26780	CDS	236	CGCGCCAGGAGAGTCCAT	0	73
	26781	CDS	258	TTAGCGGCGGGTTAGTCT	22	74
	26782	CDS	262	AGCTTTAGCGGCGGGTTA	9	75
10	26783	CDS	401	CTCGGTCTGCTTCGGGCT	30	76
	26784	CDS	409	TGCCCACACTCGGTCTGC	67	77
	26785	CDS	412	CGGTGCCCACACTCGGTC	70	78
	26786	CDS	416	GAAGCGGTGCCCACACTC	66	79
	26787	CDS	712	TTACACGCCTTCTCCACG	77	80
15	26788	CDS	716	GTATTTACACGCCTTCTC	25	81
	26789	CDS	719	CCGGTATTTACACGCCTT	35	82
	26790	CDS	816	GAGGGCAGGACACCACCA	81	83
	26791	CDS	819	TGTGAGGGCAGGACACCA	70	84
	26792	CDS	823	CACTTGAGGGCAGGAC	69	85
20	26793	CDS	939	GCTGGTTTGTCCCCCTGAA	42	86
	26794	CDS	943	ATCTGCTGTTTGTCCCC	73	87
	26795	CDS	1281	CGCGGTTCTGGAGGGACT	39	88
	26796	CDS	1316	CCCCGCACTCTTGTCAC	36	89
	26797	CDS	1319	TTGCCCCGCACTCTTGTC	19	90
25	26798	CDS	1323	CCACTTGCCCCGCACTCT	70	91
	26799	CDS	1326	GAGCCACTTGCCCCGCAC	53	92
	26800	CDS	1330	TTCCGAGCCACTTGCCCC	22	93
	26801	CDS	1485	TCCGCCGCTTGTAAGTCGC	70	94
	26802	CDS	1489	TGCTTCCGCCGCTTGTAAG	42	95
30	26803	CDS	1492	TCCTGCTTCCGCCGCTTG	73	96
	26804	CDS	1589	GTCCCCGTTTCAGGTAGAC	43	97
	26805	CDS	1593	TCCCGTCCCCGTTTCAGGT	78	98

WO 00/20435

PCT/US99/23171

-84-

	26806	CDS	1596	CCATCCCGTCCCCGTTCA	81	99
	26807	CDS	1599	TCCCCATCCCGTCCCCGT	86	100
	26808	CDS	1603	CCCTTCCCCATCCCGTCC	40	101
	26809	CDS	1609	TGCGTCCCCTTCCCCATC	45	102
5	26810	CDS	1612	AAGTGCGTCCCCTTCCCC	32	103
	26811	CDS	1616	CGACAAGTGCGTCCCCTT	71	104
	26812	CDS	1662	AAGGAAGCAGGGCATCAT	44	105
	26813	CDS	1781	CTCTCCAGTGGGCTTCTT	70	106
	26814	CDS	1785	TCATCTCTCCAGTGGGCT	48	107
10	26815	3'UTR	1933	GCTAAATCCACCTCCCCA	0	108
	26816	3'UTR	2027	TCTGCCGCTTCCTCCGTC	41	109
	26817	3'UTR	2033	CCGCCTTCTGCCGCTTCC	70	110

15 ¹Position of first nucleotide of the target site on GenBank
accession number HSU21092, incorporated herein as SEQ ID
NO: 3.

Table 14

Inhibition of TRAF-3 mRNA levels by chimeric
phosphorothioate oligonucleotides having 2'-MOE wings and a
deoxy gap

5	ISIS#	TARGET REGION	SITE ¹	SEQUENCE 5'--> 3'	%Inhib	SEQ ID NO.
	26818	5'UTR	78	AGAGCCGACGACCGCCGC	2	71
	26819	5'UTR	81	GGAAGAGCCGACGACCGC	0	72
	26820	CDS	236	CGCGCCAGGAGAGTCCAT	87	73
10	26821	CDS	258	TTAGCGGCGGGTTAGTCT	32	74
	26822	CDS	262	AGCTTTAGCGGCGGGTTA	33	75
	26823	CDS	401	CTCGGTCTGCTTCGGGCT	58	76
	26824	CDS	409	TGCCACACTCGGTCTGC	71	77
	26825	CDS	412	CGGTGCCACACTCGGTC	69	78
15	26826	CDS	416	GAAGCGGTGCCACACTC	54	79
	26827	CDS	712	TTACACGCCTTCTCCACG	65	80
	26828	CDS	716	GTATTTACACGCCTTCTC	40	81
	26829	CDS	719	CCGGTATTTACACGCCTT	77	82
	26830	CDS	816	GAGGGCAGGACACCACCA	76	83
20	26831	CDS	819	TGTGAGGGCAGGACACCA	78	84
	26832	CDS	823	CACTTGTGAGGGCAGGAC	88	85
	26833	CDS	939	GCTGGTTTGTCCCCTGAA	79	86
	26834	CDS	943	ATCTGCTGGTTTGTCCCC	74	87

WO 00/20435

PCT/US99/23171

- 86 -

	26835	CDS	1281	CGCGGTTCTGGAGGGACT	57	88
	26836	CDS	1316	CCCCGCACTCTTGTCAC	0	89
	26837	CDS	1319	TTGCCCCGCACTCTTGTC	34	90
	26838	CDS	1323	CCACTTGCCCCGCACTCT	33	91
5	26839	CDS	1326	GAGCCACTTGCCCCGCAC	39	92
	26840	CDS	1330	TTCCGAGCCACTTGCCCC	0	93
	26841	CDS	1485	TCCGCCGCTTGTAAGTCGC	71	94
	26842	CDS	1489	TGCTTCCGCCGCTTGTAAG	39	95
	26843	CDS	1492	TCCTGCTTCCGCCGCTTG	47	96
10	26844	CDS	1589	GTCCCCGTTTCAGGTAGAC	7	97
	26845	CDS	1593	TCCCGTCCCCGTTTCAGGT	56	98
	26846	CDS	1596	CCATCCCGTCCCCGTTCA	54	99
	26847	CDS	1599	TCCCCATCCCGTCCCCGT	41	100
	26848	CDS	1603	CCCTTCCCCATCCCGTCC	79	101
15	26849	CDS	1609	TGCGTCCCCCTTCCCCATC	63	102
	26850	CDS	1612	AAGTGCGTCCCCCTTCCCC	77	103
	26851	CDS	1616	CGACAAGTGCGTCCCCCTT	80	104
	26852	CDS	1662	AAGGAAGCAGGGCATCAT	4	105
	26853	CDS	1781	CTCTCCAGTGGGCTTCTT	64	106
20	26854	CDS	1785	TCATCTCTCCAGTGGGCT	55	107
	26855	3'UTR	1933	GCTAAATCCACCTCCCCA	48	108
	26856	3'UTR	2027	TCTGCCGCTTCCTCCGTC	39	109

-87-

26857 3'UTR 2033 CCGCCTTCTGCCGCTTCC 65 110

5 'Position of first nucleotide of the target site on GenBank
accession number HSU21092, incorporated herein as SEQ ID NO:
3.

Example 32

10 Antisense inhibition of TRAF-4 expression

The antisense oligonucleotides shown in Tables 7 and 8 were screened in T-24 cells for ability to inhibit human TRAF-4 expression. Results are shown in Tables 15 and 16, respectively.

15

Table 15

Inhibition of TRAF-4 mRNA levels by phosphorothioate
oligodeoxynucleotides

	ISIS#	TARGET	SITE ¹	SEQUENCE	%Inhib	SEQ ID
5	26860	Start	72	GCATGGCGGGCGAGCGGC	0	111
	26861	CDS	113	CCGTCGCTTGGGCTTCTC	16	112
	26862	CDS	232	GGGCACTTGAAGACTCCT	0	113
	26863	CDS	236	CTCAGGGCACTTGAAGAC	0	114
	26864	CDS	241	TGGTCCTCAGGGCACTTG	47	115
10	26865	CDS	245	AAGCTGGTCCTCAGGGCA	0	116
	26866	CDS	341	GCGGCAGCCCTCCTCACT	6	117
	26867	CDS	346	CTCCAGCGGCAGCCCTCC	61	118
	26868	CDS	411	TAGGGCAGGGAATGACAT	0	119
	26869	CDS	415	CGATTAGGGCAGGGAATG	20	120
15	26870	CDS	421	GGGCAGCGATTAGGGCAG	54	121
	26871	CDS	523	GCCTCCCCACTGAAGTCA	38	122
	26872	CDS	592	ATGCGGGCACCACACTTA	56	123
	26873	CDS	733	GGGCAGGCAACAGGCAGC	53	124
	26874	CDS	759	CCACAGTGCCCACACCAC	34	125
20	26875	CDS	763	CGAGCCACAGTGCCCACA	43	126
	26876	CDS	769	TCCTCCCGAGCCACAGTG	66	127
	26877	CDS	773	CAGGTCCTCCCGAGCCAC	35	128
	26878	CDS	819	GGCAGAGCACCAGGGCGG	54	129

WO 00/20435

PCT/US99/23171

- 89 -

	26879	CDS	828	CTTTGAATGGGCAGAGCA	48	130
	26880	CDS	833	GGAGTCTTTGAATGGGCA	0	131
	26881	CDS	875	ATGCCGTGCCATTGCCAG	44	132
	26882	CDS	925	CTCACCAGGGCACACATC	69	133
5	26883	CDS	944	CAGCTCCTGCCGTTGCCG	71	134
	26884	CDS	1000	ATGAGCACGCCATCACTG	0	135
	26885	CDS	1033	TGTAGCCGCCGTCCATAG	74	136
	26886	CDS	1039	GCCTCCTGTAGCCGCCGT	53	137
	26887	CDS	1081	TAGAAGGCTGGGCTGAAG	48	138
10	26888	CDS	1086	GTGTGTAGAAGGCTGGGC	23	139
	26889	CDS	1152	GTGTGCCCTCACCCTGC	32	140
	26890	CDS	1226	GACACGGCGGGCAAAGGG	52	141
	26891	CDS	1232	GAAGGTGACACGGCGGGC	73	142
	26892	CDS	1260	GCCCAGGGTCGCTCTGAT	80	143
15	26893	CDS	1313	CTTCCAGTTTGGGTCGGG	0	144
	26894	CDS	1377	GATAACCAAAGCCCAGAG	45	145
	26895	3'UTR	1513	CATCGTCCTTTCCCCTCG	51	146
	26896	3'UTR	1660	GGCCAGGGCTGAAGCACC	53	147
	26897	3'UTR	1703	TTGTTTCCAGCCCTTCAT	67	148
20	26898	3'UTR	1746	CATGTCTGCCCTACCCAA	0	149
	26899	3'UTR	1948	GCTCCCCTGCTGTGCCCT	49	150

WO 00/20435

PCT/US99/23171

-90-

¹Position of first nucleotide of the target site on GenBank
Accession No. X80200, locus name "HSMLN62" SEQ ID NO. 4.

-91-

Table 16

Inhibition of TPAF-4 mRNA levels by chimeric
phosphorothioate oligonucleotides having 2'-MOE wings and a
deoxy gap

5						
	ISIS#	TARGET	SITE ¹	SEQUENCE	%Inhib	SEQ ID
	26901	CDS	113	CCCTCGCTTGGGCTTCTC	25	112
	26902	CDS	232	GGGCACTTGAAGACTCCT	57	113
	26903	CDS	236	CTCAGGGCACTTGAAGAC	49	114
10	26904	CDS	241	TGGTCCTCAGGGCACTTG	0	115
	26905	CDS	245	AAGCTGGTCCTCAGGGCA	0	116
	26906	CDS	341	GCGGCAGCCCTCCTCACT	0	117
	26907	CDS	346	CTCCAGCGGCAGCCCTCC	68	118
	26908	CDS	411	TAGGGCAGGGAATGACAT	0	119
15	26909	CDS	415	CGATTAGGGCAGGGAATG	0	120
	26910	CDS	421	GGGCAGCGATTAGGGCAG	39	121
	26911	CDS	523	GCCTCCCCACTGAAGTCA	30	122
	26912	CDS	592	ATGCGGGCACCACACTTA	44	123
	26913	CDS	733	GGGCAGGCAACAGGCAGC	66	124
20	26914	CDS	759	CCACAGTGCCACACCAC	43	125
	26915	CDS	763	CGAGCCACAGTGCCACAC	6	126
	26916	CDS	769	TCCTCCCGAGCCACAGTG	46	127
	26917	CDS	773	CAGGTCCTCCCGAGCCAC	75	128

	ISIS#	TARGET	SITE ¹	SEQUENCE	%Inhib	SEQ ID
	26918	CDS	819	GGCAGAGCACCAGGGCGG	74	129
	26919	CDS	828	CTTTGAATGGGCAGAGCA	34	130
	26920	CDS	833	GGAGTCTTTGAATGGGCA	11	131
	26921	CDS	875	ATGCCGTGCCATTGCCAG	32	132
5	26922	CDS	925	CTCACCAGGGCACACATC	30	133
	26923	CDS	944	CAGCTCCTGCCGTTGCCG	79	134
	26924	CDS	1000	ATGAGCACGCCATCACTG	39	135
	26925	CDS	1033	TGTAGCCGCCGTCCATAG	24	136
	26926	CDS	1039	GCCTCCTGTAGCCGCCGT	61	137
10	26927	CDS	1081	TAGAAGGCTGGGCTGAAG	51	138
	26928	CDS	1086	GTGTGTAGAAGGCTGGGC	75	139
	26929	CDS	1152	GTGTGCCCTCACCCTGC	23	140
	26930	CDS	1226	GACACGGCGGGCAAAGGG	27	141
	26931	CDS	1232	GAAGGTGACACGGCGGGC	65	142
15	26932	CDS	1260	GCCCAGGGTCGCTCTGAT	76	143
	26933	CDS	1313	CTTCCAGTTTGGGTCGGG	0	144
	26934	CDS	1377	GATAACCAAAGCCCAGAG	0	145
	26935	3'UTR	1513	CATCGTCCTTTCCCCTCG	11	146
	26936	3'UTR	1660	GGCCAGGGCTGAAGCACC	79	147
20	26937	3'UTR	1703	TTGTTTCCAGCCCTTCAT	7	148

-93-

ISIS#	TARGET	SITE ¹	SEQUENCE	%Inhib	SEQ ID
26938	3'UTR	1746	CATGTCTGCCCTACCCAA	26	149
26939	3'UTR	1948	GCTCCCCTGCTGTGCCCT	14	150

¹Position of first nucleotide of the target site on GenBank
5 Accession No. X80200, locus name "HSMLN62" SEQ ID NO. 4.

Example 32

Antisense inhibition of TRAF-5 expression

The antisense oligonucleotides shown in Tables 9 and 10
10 were screened in T-24 cells for ability to inhibit human
TRAF-5 expression. Results are shown in Tables 17 and 18,
respectively.

Table 17

Inhibition of TRAF-5 mRNA levels by phosphorothioate
oligodeoxynucleotides

	ISIS#	TARGET	SITE ¹	SEQUENCE	%Inhib	SEQ ID
5	26940	Start	49	TGAATAAGCCATTGTGGG	33	151
	26941	CDS	62	CTTTATGCTCTTCTGAAT	0	152
	26942	CDS	83	GGATGAAACCACAGGGCA	14	153
	26943	CDS	120	TCAAAGTCCAAGGAAATG	37	154
	26944	CDS	195	TGAAGCACCGAGTGGCAG	37	155
10	26945	CDS	282	GGGCAGATTGGCACTGTG	79	156
	26946	CDS	313	CTCCTGAGATTTGATGAC	0	157
	26947	CDS	491	CTTTCCGTAGGACTGGCT	60	158
	26948	CDS	584	GATTCTGTAGATTGATGA	2	159
	26949	CDS	667	TTCATCTACCTCAGTTTT	55	160
15	26950	CDS	735	TCCGTTACAGCACAGCCA	59	161
	26951	CDS	788	GCATGTGCTCCCGTAAGG	77	162
	26952	CDS	907	CTTTTCAAGTTTCTTTAT	4	163
	26953	CDS	1079	CTTCCATCAAAGGTCTCA	35	164
	26954	CDS	1146	TCTAAAACGGCTAATCTT	0	165
20	26955	CDS	1283	TCATCTTGTAATCTGTCA	9	166
	26956	CDS	1333	GGACTGGCTGAAGATGGA	7	167
	26957	CDS	1403	CCCTCCCTGACCCATCCC	71	168

-95-

	ISIS#	TARGET	SITE ¹	SEQUENCE	%Inhib	SEQ ID
	26958	CDS	1620	GAATGAGCCACAAAGCGG	68	169
	26959	CDS	1672	CAAGAACAGAGTGTTCATC	13	170
	26960	3'UTR	1799	GTCTAAATCCAGGTCAAT	26	171
	26961	3'UTR	1964	AAACTTACCATCTTTCAA	48	172
5	26962	3'UTR	2053	CTCTGTGTCTCCATAAC	54	173
	26963	3'UTR	2167	CTTAAGTGAACAGCCTA	35	174
	26964	3'UTR	2352	GCAGGAAGAATGAAAATG	0	175
	26965	3'UTR	2501	TATTTGGTTGAATCTTAT	8	176
	26966	3'UTR	2611	AAATTCTATCCATCCTCA	32	177
10	26967	3'UTR	2683	AAATTGTAAAGGTTTTCT	22	178
	26968	3'UTR	2779	ACAATGAACTCTGTCTC	14	179
	26969	3'UTR	2940	GCAAACTCCGTCTCTAC	51	180
	26970	3'UTR	3055	CAATAGTTGTCAGAGGCT	39	181
	26971	3'UTR	3209	AAGGACTCATCTCAGTTT	0	182
15	26972	3'UTR	3280	TAAACAACGCAGAAGGGCT	74	183
	26973	3'UTR	3295	AGTAGGGAAGTGGCATAA	29	184
	26974	3'UTR	3377	CATCACCAGGTAAGCAGC	60	185
	26975	3'UTR	3553	TCCTGTTGTGAACCTATT	40	186
	26976	3'UTR	3656	GGAATTGTGGGCTAAAGA	60	187
20	26977	3'UTR	3724	GCTCAGGAAGACAGAGTG	6	188

WO 00/20435

PCT/US99/23171

-96-

ISIS#	TARGET	SITE ¹	SEQUENCE	%Inhib	SEQ ID
26978	3'UTR	3873	TGAACTCCTAAGCAAACC	23	189
26979	3'UTR	3889	GATGATGAAGGAACTCTG	20	190

- 5 ¹Position of first nucleotide of the target site on GenBank
GenBank Accession No. AB000509, locus name "AB000509" SEQ
ID NO. 5

-97-

Table 18

Inhibition of TRAF-5 mRNA levels by chimeric
phosphorothioate oligonucleotides having 2'-MOE wings and a
deoxy gap

5	ISIS#	TARGET	SITE ¹	SEQUENCE	%Inhib	SEQ ID
	26980	Start	49	TGAAT ¹ AGCCATTGTGGG	27	151
	26981	CDS	62	CTTTATGCTCTTCTGAAT	43	152
	26982	CDS	83	GGATGAAACCACAGGGCA	71	153
10	26983	CDS	120	TCAAAGTCCAAGGAAATG	48	154
	26984	CDS	195	TGAAGCACCGAGTGGCAG	66	155
	26985	CDS	282	GGGCAGATTGGCACTGTG	37	156
	26986	CDS	313	CTCCTGAGATTTGATGAC	64	157
	26987	CDS	491	CTTTCCGTAGGACTGGCT	71	158
15	26988	CDS	584	GATTCTGTAGATTGATGA	18	159
	26989	CDS	667	TTCATCTACCTCAGTTTT	50	160
	26990	CDS	735	TCCGTTACAGCACAGCCA	66	161
	26991	CDS	788	GCATGTGCTCCCGTAAGG	83	162
	26992	CDS	907	CTTTTCAAGTTTCTTTAT	39	163
20	26993	CDS	1079	CTTCCATCAAAGGTCTCA	82	164
	26994	CDS	1146	TCTAAAACGGCTAATCTT	38	165
	26995	CDS	1283	TCATCTTGTAATCTGTCA	61	166
	26996	CDS	1333	GGACTGGCTGAAGATGGA	35	167

- 98 -

	ISIS#	TARGET	SITE ¹	SEQUENCE	%Inhib	SEQ ID
	26997	CDS	1403	CCCTCCCTGACCCATCCC	40	168
	26998	CDS	1620	GAATGAGCCACAAAGCGG	76	169
	26999	CDS	1672	CAAGAACAGAGTGTCATC	25	170
	27000	3'UTR	1799	GTCTAAATCCAGGTCAAT	30	171
5	27001	3'UTR	1964	AAACTTACCATCTTTCAA	66	172
	27002	3'UTR	2053	CTCTGTGTCCTCCATAAC	68	173
	27003	3'UTR	2167	CTTAACTGGAACAGCCTA	68	174
	27004	3'UTR	2352	GCAGGAAGAATGAAAATG	20	175
	27005	3'UTR	2501	TATTTGGTTGAATCTTAT	38	176
10	27006	3'UTR	2611	AAATTCTATCCATCCTCA	0	177
	27007	3'UTR	2683	AAATTGTAAAGGTTTCT	8	178
	27008	3'UTR	2779	ACAATGAAACTCTGTCTC	66	179
	27009	3'UTR	2940	GCAAAACTCCGTCTCTAC	51	180
	27010	3'UTR	3055	CAATAGTTGTCAGAGGCT	32	181
15	27011	3'UTR	3209	AAGGACTCATCTCAGTTT	20	182
	27012	3'UTR	3280	TAACAACGCAGAAGGGCT	64	183
	27013	3'UTR	3295	AGTAGGGAAGTGGCATAA	58	184
	27014	3'UTR	3377	CATCACCAGGTAAGCAGC	59	185
	27015	3'UTR	3553	TCCTGTTGTGAACCTATT	79	186
20	27016	3'UTR	3656	GGACTTGTGGGCTAAAGA	67	187

-99-

ISIS#	TARGET	SITE ¹	SEQUENCE	%Inhib	SEQ ID
27017	3'UTR	3724	GCTCAGGAAGACAGAGTG	62	188
27018	3'UTR	3873	TGAACTCCTAAGCAAACC	16	189
27019	3'UTR	3889	GATGATGAAGGAACTCTG	52	190

- 5 ¹Position of first nucleotide of the target site on GenBank Accession No. AB000509, locus name "AB000509" SEQ ID NO. 5.

Example 33**Additional oligonucleotides targeted to TRAF-1**

- 10 Additional antisense oligonucleotides targeted to TRAF-1 were designed and synthesized. All compounds are 2'-MOE gapmers with 2'-MOE nucleotides shown in bold. Backbones are phosphorothioates throughout. All C are 5-methyl C. Compounds were tested as described in Example 13 above. Compounds and
- 15 results are shown in Table 19.

Table 19**Antisense inhibition of TRAF-1 expression**

ISIS #	SEQUENCE	GENE TARGET REGION	START POS ¹	% INHIB	SEQ ID NO:
20 101869	GGACCAGCCTTGTGGAGTCC	5' UTR	5	--	229
101870	TCAGGGTTCCAGGCTGGCCA	5' UTR	55	--	230
101871	TCTCAGGGTTCCAGGCTGGC	Start	57	51	231
101872	CATCTCAGGGTTCCAGGCTG	Start	59	50	232
101873	GCCATCTCAGGGTTCCAGGC	Start	61	5	233
25 101874	AGGCCATCTCAGGGTTCCAG	Start	63	--	234
101875	GGAGGCCATCTCAGGGTTCC	Start	65	--	235
101876	CTGGAGGCCATCTCAGGGTT	Start	67	7	236
101877	AGCTGGAGGCCATCTCAGGG	Start	69	--	237

-100-

ISIS #	SEQUENCE	GENE TARGET REGION	START POS ¹	% INHIB	SEQ ID NO:
101878	TGAGCTGGAGGCCATCTCAG	Start	71	--	238
101879	CCTGAGCTGGAGGCCATCTC	Start	73	28	239
101880	TGCCTGAGCTGGAGGCCATC	Start	75	--	240
101881	GCTGCCTGAGCTGGAGGCCA	Start	77	--	241
5 101882	GGGCGAGGACTGCTGCCTGA	Coding	88	60	242
101883	TCTCAGAGAGACAGCCTGCA	Coding	189	43	243
101884	TCCTGGGCTTATAGACTGGA	Coding	260	78	244
101885	GGGCTTCCCTTGAAGGAGCA	Coding	358	--	245
101886	CAACAGCAGGTTTAGGTGGG	Coding	416	--	246
10 101887	CAGGGCCATGGGCCAGACT	Coding	479	--	247
101888	CCCGGTAGCAATCGACCTCC	Coding	555	31	248
101889	CTCAGCCAGAAGCTTCTCCT	Coding	623	53	249
101890	GGCCAGGGCCAGGTGGGAGG	Coding	704	--	250
101891	TGCTGAAGCTCCACCACCCT	Coding	778	11	251
15 101892	GTGATCTTCCACAGGAAAGT	Coding	877	8	252
101893	TTGGCAGTGTTAGAAGGCTGG	Coding	955	27	253
101894	AAGAGCGACAGATGGGTTCT	Coding	1030	61	254
101895	GCAGCATGAAGGTGACCTTG	Coding	1104	--	255
101896	TCTGGAAGGACGCTGAGCTT	Coding	1173	--	256
20 101897	CTCCACAATGCACTTGAGGA	Coding	1295	35	257
101898	TGCCCTGAGGTCTTGGGTGC	3' UTR	1399	31	258
101899	ATCCTAACCAGATGGCCAGC	3' UTR	1473	10	259
101900	ACTGGCCTCCCAGTGTGCGCA	3' UTR	1571	9	260
101901	GCTTGGGTCTTACGGTTCCA	3' UTR	1656	22	261
25 101902	TCCTGTTTCTGACCCTGGAG	3' UTR	1749	81	262
101903	CTGGGTTTGTCTGTTTCACCT	3' UTR	1826	85	263
101904	CCAGGAGGCTAGAATGAGAG	3' UTR	1917	--	264
101905	TGAGGAGCTGGGAGGACAGG	3' UTR	1986	--	265
101906	TTGGGAAGCTGAGCTGCCAG	3' UTR	2066	--	266

-101-

ISIS #	SEQUENCE	GENE	START	%	SEQ ID
		TARGET	POS ¹	INHIB	NO:
		REGION			
101907	ACAGTGACTGGGTTTCACCT	3' UTR	2153	35	267
101908	TTACGGGATTCTGGAAAGCA	3' UTR	2256	78	268

¹Position of first nucleotide of the target site on GenBank
5 Accession No.U19261, locus name "HSU19261," SEQ ID NO: 1.

-102-

What is claimed is:

1. An antisense compound 8 to 30 nucleotides in length targeted to a nucleic acid molecule encoding a human tumor
5 necrosis factor receptor-associated factor, wherein said antisense compound inhibits the expression of human tumor necrosis factor receptor-associated factor.
2. The antisense compound of claim 1 which is an
10 antisense oligonucleotide.
3. The antisense oligonucleotide of claim 2 which comprises at least one modified internucleoside linkage.
- 15 4. The antisense oligonucleotide of claim 3 wherein the modified internucleoside linkage is a phosphorothioate linkage.
5. The antisense oligonucleotide of claim 2 which
20 comprises at least one modified sugar moiety.
6. The antisense oligonucleotide of claim 5 wherein the modified sugar moiety is a 2'-O-methoxyethyl sugar moiety.
- 25 7. The antisense oligonucleotide of claim 2 which comprises at least one modified nucleobase.
8. The antisense oligonucleotide of claim 7 wherein the modified nucleobase is a 5-methylcytosine.
30
9. The antisense compound of claim 2 which is a chimeric oligonucleotide.

-103-

10. The antisense compound of claim 1 wherein the human TRAF is TRAF-2 or TRAF-6.

11. A composition comprising the antisense compound of
5 claim 1 and a pharmaceutically acceptable carrier or diluent.

12. The composition of claim 11 comprising a colloidal dispersion system.

10 13. The composition of claim 11 wherein the antisense compound is an antisense oligonucleotide.

14. A method of inhibiting the expression of tumor necrosis factor receptor-associated factor in human cells or
15 tissues comprising contacting said cells or tissues with the antisense compound of claim 1 so that expression of tumor necrosis factor receptor-associated factor is inhibited.

15. A method of treating a human having a disease or
20 condition associated with tumor necrosis factor receptor-associated factor comprising administering to said animal a therapeutically or prophylactically effective amount of the antisense compound of claim 1 so that expression of tumor necrosis factor receptor-associated factor is inhibited.

25 16. The method of claim 15 wherein the disease or condition is a hyperproliferative or inflammatory disease or condition.

30 17. A method of reducing jun kinase activation in cells or tissues by tumor necrosis factor- α comprising contacting said cells or tissues with an antisense compound targeted to TRAF-2.

35 18. A method of reducing jun kinase activation in cells

-104-

or tissues comprising contacting said cells or tissues with an antisense compound targeted to TRAF-6.

19. A method of reducing E-selectin expression in cells
5 or tissues comprising contacting said cells or tissues with an antisense compound targeted to TRAF-2 or TRAF-6.

SEQUENCE LISTING

<110> Baker, Brenda F.
Cowser, Lex M.
Monia, Brett P.
Xu, Xiaoxing S.
Isis Pharmaceuticals, Inc.

<120> ANTISENSE MODULATION OF EXPRESSION OF TUMOR NECROSIS FACTOR
RECEPTOR-ASSOCIATED FACTOR (TRAFs)

<130> ISPH-0411

<140> 09/167,109

<141> 1998-10-06

<160> 268

<210> 1

<211> 2380

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (76)..(1326)

<300>

<308> U19261 Genbank

<309> 1995-02-21

<400> 1

gccaggactc cacaaggctg gtccctgcc ctggagcaac ttaaacaggc cctctggcca 60

gcctggaacc ctgag atg gcc tcc agc tca ggc agc agt cct cgc ccg gcc 111
Met Ala Ser Ser Ser Gly Ser Ser Pro Arg Pro Ala
1 5 10

cct gac gag aat gag ttt ccc ttt ggg tgc cct ccc acc gtc tgc cag 159
Pro Asp Glu Asn Glu Phe Pro Phe Gly Cys Pro Pro Thr Val Cys Gln
15 20 25

gac cca aag gag ccc agg gct ctc tgc tgt gca ggc tgt ctc tct gag 207
Asp Pro Lys Glu Pro Arg Ala Leu Cys Cys Ala Gly Cys Leu Ser Glu
30 35 40

aac ccg agg aat ggc gag gat cag atc tgc ccc aaa tgc aga ggg gaa 255
Asn Pro Arg Asn Gly Glu Asp Gln Ile Cys Pro Lys Cys Arg Gly Glu
45 50 55 60

gac ctc cag tct ata agc cca gga agc cgt ctt cga act cag gag aag 303
Asp Leu Gln Ser Ile Ser Pro Gly Ser Arg Leu Arg Thr Gln Glu Lys
65 70 75

gct cac ccc gag gtg gct gag gct gga att ggg tgc ccc ttt gca ggt 351

Ala His Pro Glu Val Ala Glu Ala Gly Ile Gly Cys Pro Phe Ala Gly
80 85 90

gtc ggc tgc tcc ttc aag gga agc cca cag tct gtg caa gag cat gag 399
Val Gly Cys Ser Phe Lys Gly Ser Pro Gln Ser Val Gln Glu His Glu
95 100 105

WO 00/20435

PCT/US99/23171

gtc acc tcc cag acc tcc cac cta aac ctg ctg ttg ggg ttc atg aaa	447
Val Thr Ser Gln Thr Ser His Leu Asn Leu Leu Leu Gly Phe Met Lys	
110 115 120	
cag tgg aag gcc cgg ctg ggc tgt ggc ctg gag tct ggg ccc atg gcc	495
Gln Trp Lys Ala Arg Leu Gly Cys Gly Leu Glu Ser Gly Pro Met Ala	
125 130 135 140	
ctg gag cag aac ctg tca gac ctg cag ctg cag gca gcc gtg gaa gtg	543
Leu Glu Gln Asn Leu Ser Asp Leu Gln Leu Gln Ala Ala Val Glu Val	
145 150 155	
gcg ggg gac ctg gag gtc gat tgc tac cgg gca ccc tgc tcc gag agc	591
Ala Gly Asp Leu Glu Val Asp Cys Tyr Arg Ala Pro Cys Ser Glu Ser	
160 165 170	
cag gag gag ctg gcc ctg cag cac ttc atg aag gag aag ctt ctg gct	639
Gln Glu Glu Leu Ala Leu Gln His Phe Met Lys Glu Lys Leu Leu Ala	
175 180 185	
gag ctg gag ggg aag ctg cgt gtg ttt gag aac att gtt gct gtc ctc	687
Glu Leu Glu Gly Lys Leu Arg Val Phe Glu Asn Ile Val Ala Val Leu	
190 195 200	
aac aag gag gtg gag gcc tcc cac ctg gcc ctg gcc acc tct atc cac	735
Asn Lys Glu Val Glu Ala Ser His Leu Ala Leu Ala Thr Ser Ile His	
205 210 215 220	
cag agc cag ctg gac cgt gag cgc atc ctg agc ttg gag cag agg gtg	783
Gln Ser Gln Leu Asp Arg Glu Arg Ile Leu Ser Leu Glu Gln Arg Val	
225 230 235	
gtg gag ctt cag cag acc ctg gcc cag aaa gac cag gcc ctg ggc aag	831
Val Glu Leu Gln Gln Thr Leu Ala Gln Lys Asp Gln Ala Leu Gly Lys	
240 245 250	
ctg gag cag agc ttg cgc ctc atg gag gag gcc tcc ttc gat ggc act	879
Leu Glu Gln Ser Leu Arg Leu Met Glu Glu Ala Ser Phe Asp Gly Thr	
255 260 265	
ttc ctg tgg aag atc acc aat gtc acc agg cgg tgc cat gag tcg gcc	927
Phe Leu Trp Lys Ile Thr Asn Val Thr Arg Arg Cys His Glu Ser Ala	
270 275 280	
tgt ggc agg acc gtc agc ctc ttc tcc cca gcc ttc tac act gcc aag	975
Cys Gly Arg Thr Val Ser Leu Phe Ser Pro Ala Phe Tyr Thr Ala Lys	
285 290 295 300	
tat ggc tac aag ttg tgc ctg cgg ctg tac ctg aat gga gat ggc act	1023
Tyr Gly Tyr Lys Leu Cys Leu Arg Leu Tyr Leu Asn Gly Asp Gly Thr	
305 310 315	
gga aag aga acc cat ctg tgc ctc ttc atc gtg atc atg aga ggg gag	1071
Gly Lys Arg Thr His Leu Ser Leu Phe Ile Val Ile Met Arg Gly Glu	
320 325 330	
tat gat gcg ctg ctg ccg tgg ccc ttc cgg aac aag gtc acc ttc atg	1119
Tyr Asp Ala Leu Leu Pro Trp Pro Phe Arg Asn Lys Val Thr Phe Met	
335 340 345	
ctg ctg gac cag aac aac cgt gag cac gcc att gac gcc ttc cgg cct	1167
Leu Leu Asp Gln Asn Asn Arg Glu His Ala Ile Asp Ala Phe Arg Pro	
350 355 360	


```

gac cta agc tca gcg tcc ttc cag agg ccc cag agt gaa acc aac gtg 1215
Asp Leu Ser Ser Ala Ser Phe Gln Arg Pro Gln Ser Glu Thr Asn Val
365 370 375 380

gcc agt gga tgc cca ctc ttc ttc ccc ctc agc aaa ctg cag tca ccc 1263
Ala Ser Gly Cys Pro Leu Phe Phe Pro Leu Ser Lys Leu Gln Ser Pro
385 390 395

aag cac gcc tac gtg aag gac gac aca atg ttc ctc aag tgc att gtg 1311
Lys His Ala Tyr Val Lys Asp Asp Thr Met Phe Leu Lys Cys Ile Val
400 405 410

gag acc agc act tag ggtggggcggg gctcctgagg gagctccaac tcagaaggga 1366
Glu Thr Ser Thr
415

gctagccaga ggactgtgat gccctggcct tggcacccaa gacctcaggg cacaaagatg 1426
ggtgaaggct ggcatgatcc aagcaagact gaggggtcga cttcgggctg gccatctggt 1486
taggatggca ggacgtgggc tgggcccaca aaggcaaagg gtccagaagg agacaggcag 1546
agctgctccc ctctgcacgg accatgogac actgggaggg cagtgaagcca ctccggcccc 1606
gaatgttgag gtggactctc accaaatgag aagaaaatgg aaccaggctt ggaaccgtag 1666
gacccaagca gagaagctct cgggctagga agatctctgc agggccgcca gggagacctg 1726
gacacaggcc tgctctcttt ttctccaggg tcagaaacag gaccgggtgg aagggatggg 1786
gtgccagttt gaatgcagtc tgtccaggct cgtcattgga ggtgaacaag caaacccaga 1846
cggctccact aggacttcaa attggggggt ggatttgaag acttttaagt ttcttccag 1906
cccagaaagt ctctcattct agcctcctgg ccagggtgag tcttagagct acaggggttc 1966
tggaacatt caggagcttc ctgtcctccc agctcctcac tcaccttcag taacccccac 2026
tggactgacc tggccacag ggcacctgcc accctgggcc tggcagctca gcttcccaac 2086
acgcaggagc acaccagcc cccacatcct gtgcctccat cagctaaaca ccacgtcact 2146
tcatgcaggt gaaacccagt cactgtgagc tcccaggtgc agccagaggc acctcaagaa 2206
gaagaggggc ataaactttc ctcttctgct ctagaggccc cacctttggt gctttccaga 2266
atcccgtaac acctgattaa ctgaggcatc cacttctttc agcagactga tcaggacctc 2326
caagccactg agcaatgtat aacccccaaag ggaattcaaa aaaaaaaaaa 2380

<210> 2
<211> 2262
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (55)..(1560)

<300>
<308> U12597 Genbank
<309> 1996-02-16

```

<400> 2

```

gaattccggc gcgctgcgac cggtggggct ttgttcgcgg gggtcacagc tctc atg 57
                                     Met
                                     1

gct gca gct agc gtg acc ccc cct ggc tcc ctg gag ttg cta cag ccc 105
Ala Ala Ala Ser Val Thr Pro Pro Gly Ser Leu Glu Leu Leu Gln Pro
      5                      10                      15

ggc ttc tcc aag acc ctc ctg ggg acc aag ctg gaa gcc aag tac ctg 153
Gly Phe Ser Lys Thr Leu Leu Gly Thr Lys Leu Glu Ala Lys Tyr Leu
      20                      25                      30

tgc tcc gcc tgc aga aac gtc ctc cgc agg ccc ttc cag gcg cag tgt 201
Cys Ser Ala Cys Arg Asn Val Leu Arg Arg Pro Phe Gln Ala Gln Cys
      35                      40                      45

ggc cac cgg tac tgc tcc ttc tgc ctg gcc agc atc ctc agc tct ggg 249
Gly His Arg Tyr Cys Ser Phe Cys Leu Ala Ser Ile Leu Ser Ser Gly
      50                      55                      60                      65

cct cag aac tgt gct gcc tgt gtt cac gag ggc ata tat gaa gaa ggc 297
Pro Gln Asn Cys Ala Ala Cys Val His Glu Gly Ile Tyr Glu Glu Gly
      70                      75                      80

att tct att tta gaa agc agt tgc gcc ttc cca gat aat gct gcc cgc 345
Ile Ser Ile Leu Glu Ser Ser Ser Ala Phe Pro Asp Asn Ala Ala Arg
      85                      90                      95

agg gag gtg gag agc ctg ccg gcc gtc tgt ccc agt gat gga tgc acc 393
Arg Glu Val Glu Ser Leu Pro Ala Val Cys Pro Ser Asp Gly Cys Thr
      100                     105                     110

tgg aag ggg acc ctg aaa gaa tac gag agc tgc cac gaa ggc cgc tgc 441
Trp Lys Gly Thr Leu Lys Glu Tyr Glu Ser Cys His Glu Gly Arg Cys
      115                     120                     125

ccg ctc atg ctg acc gaa tgt ccc gcg tgt aaa ggc ctg gtc cgc ctt 489
Pro Leu Met Leu Thr Glu Cys Pro Ala Cys Lys Gly Leu Val Arg Leu
      130                     135                     140                     145

ggt gaa aag gag cgc cac ctg gag cac gag tgc ccg gag aga agc ctg 537
Gly Glu Lys Glu Arg His Leu Glu His Glu Cys Pro Glu Arg Ser Leu
      150                     155                     160

agc tgc cgg cat tgc cgg gca ccc tgc tgc gga gca gac gtg aag gcg 585
Ser Cys Arg His Cys Arg Ala Pro Cys Cys Gly Ala Asp Val Lys Ala
      165                     170                     175

cac cac gag gtc tgc ccc aag ttc ccc tta act tgt gac ggc tgc ggc 633
His His Glu Val Cys Pro Lys Phe Pro Leu Thr Cys Asp Gly Cys Gly
      180                     185                     190

aag aag aag atc ccc cgg gag aag ttt cag gac cac gtc aag act tgt 681
Lys Lys Lys Ile Pro Arg Glu Lys Phe Gln Asp His Val Lys Thr Cys
      195                     200                     205

ggc aag tgt cga gtc cct tgc aga ttc cac gcc atc ggc tgc ctc gag 729
Gly Lys Cys Arg Val Pro Cys Arg Phe His Ala Ile Gly Cys Leu Glu
      210                     215                     220                     225

acg gta gag ggt gag aaa cag cag gag cac gag gtg cag tgg ctg cgg 777

```

Thr Val Glu Gly Glu Lys Gln Gln Glu His Glu Val Gln Trp Leu Arg	
230 235 240	
gag cac ctg gcc atg cta ctg agc tcg gtg ctg gag gca aag ccc ctc	825
Glu His Leu Ala Met Leu Leu Ser Ser Val Leu Glu Ala Lys Pro Leu	
245 250 255	
ttg gga gac cag agc cac gcg ggg tca gag ctc ctg cag agg tgc gag	873
Leu Gly Asp Gln Ser His Ala Gly Ser Glu Leu Leu Gln Arg Cys Glu	
260 265 270	
agc ctg gag aag aag acg gcc act ttt gag aac att gtc tgc gtc ctg	921
Ser Leu Glu Lys Lys Thr Ala Thr Phe Glu Asn Ile Val Cys Val Leu	
275 280 285	
aac cgg gag gtg gag agg gtg gcc atg act gcc gag gcc tgc agc cgg	969
Asn Arg Glu Val Glu Arg Val Ala Met Thr Ala Glu Ala Cys Ser Arg	
290 295 300 305	
cag cac cgg ctg gac caa gac aag att gaa gcc ctg agt agc aag gtg	1017
Gln His Arg Leu Asp Gln Asp Lys Ile Glu Ala Leu Ser Ser Lys Val	
310 315 320	
cag cag ctg gag agg agc att ggc ctc aag gac ctg gcg atg gct gac	1065
Gln Gln Leu Glu Arg Ser Ile Gly Leu Lys Asp Leu Ala Met Ala Asp	
325 330 335	
ttg gag cag aag gtc agg ccc ttc cag gcg cag tgt ggc cac cgg tac	1113
Leu Glu Gln Lys Val Arg Pro Phe Gln Ala Gln Cys Gly His Arg Tyr	
340 345 350	
tgc tcc ttc tgc ctg gcc agc atc ctc agg aag ctc cag gaa gct gtg	1161
Cys Ser Phe Cys Leu Ala Ser Ile Leu Arg Lys Leu Gln Glu Ala Val	
355 360 365	
gct ggc cgc ata ccc gcc atc ttc tcc cca gcc ttc tac acc agc agg	1209
Ala Gly Arg Ile Pro Ala Ile Phe Ser Pro Ala Phe Tyr Thr Ser Arg	
370 375 380 385	
tac ggc tac aag atg tgt ctg cgt atc tac ctg aac ggc gac gcc acc	1257
Tyr Gly Tyr Lys Met Cys Leu Arg Ile Tyr Leu Asn Gly Asp Gly Thr	
390 395 400	
ggg cga gga aca cac ctg tcc ctc ttc ttt gtg gtg atg aag ggc cgg	1305
Gly Arg Gly Thr His Leu Ser Leu Phe Phe Val Val Met Lys Gly Pro	
405 410 415	
aat gac gcc ctg ctg cgg tgg ccc ttc aac cag aag gtg acc tta atg	1353
Asn Asp Ala Leu Leu Arg Trp Pro Phe Asn Gln Lys Val Thr Leu Met	
420 425 430	
ctg ctc gac cag aat aac cgg gag cac gtg att gac gcc ttc agg ccc	1401
Leu Leu Asp Gln Asn Asn Arg Glu His Val Ile Asp Ala Phe Arg Pro	
435 440 445	
gac gtg act tca tcc tct ttt cag agg cca gtc aac gac atg aac atc	1449
Asp Val Thr Ser Ser Ser Phe Gln Arg Pro Val Asn Asp Met Asn Ile	
450 455 460 465	
gca agc ggc tgc ccc ctc ttc tgc ccc gtc tcc aag atg gag gca aag	1497
Ala Ser Gly Cys Pro Leu Phe Cys Pro Val Ser Lys Met Glu Ala Lys	
470 475 480	

PCT/US99/23171

ctg aca ggg ctc taa ctgcccccta ctggtgtctg ggggttgggg gcagccaggc 1600
Leu Thr Gly Leu
500

acagccggct	cacggagggg	ccaccacgct	gggccagggt	ctcactgtac	aagtgggcag	1660
gggccccgct	tgggcgcttg	ggaggggtgtc	ggcctgcagc	caagttcact	gtcacggggg	1720
aaggagccac	cagccagtcc	tcagatttca	gagactgcgg	aggggcttgg	cagacggtct	1780
tagccaaggg	ctgtggtggc	attggccgag	ggtcttcggg	tgcttcccag	cacaagctgc	1840
ccttgctgtc	ctgtgcagtg	aagggagagg	ccctgggttg	gggacactca	gagtgggagc	1900
acatcccagc	agtgcccatg	tagcaggagc	acagtggatg	gccttgctgtc	cctcgggcat	1960
gacaggcaga	aacgagggct	gctccaggag	aagggcctcc	tgctggccag	agcaaggaag	2020
gctgagcagc	ttggttctcc	cctctggccc	ctggagagaa	gggagcattc	ctagacccct	2080
gggtgcttgt	ctgcacagag	ctctggtctg	tgccaccttg	gccaggctgg	ctgtgggagg	2140
gtctggtccc	acgccgcctc	tgctcagaca	ctgtgtggga	gggcacagca	cagctgcggg	2200
taaagtgtga	gagcttgcca	tccagctcac	gaagacagag	ttattaaacc	attacaaatc	2260
tc						2262

```
<210>      3
<211>      2455
<212>      DNA
<213>      Homo sapiens
```

```
<220>
<221> CDS
<222> (218) .. (1924)
```

<300>
<308> U21092
<309> 1995-03-23

<400> 3

[illegible]

atg gac tct cct ggc gcg ctg cag act aac ccg ccg cta aag ctg cac 283
Met Asp Ser Pro Gly Ala Leu Gln Thr Asn Pro Pro Leu Lys Leu His
10 15 20

act gac cgt agt gct ggg acg cca gtt ttt gtc cct gaa caa gga ggt 331
Thr Asp Arg Ser Ala Gly Thr Pro Val Phe Val Pro Glu Gln Gly Gly

25	30	35	
tac aag gaa aag ttt gtg aag acc gtg gag gac aag tac aag tgt gag Tyr Lys Glu Lys Phe Val Lys Thr Val Glu Asp Lys Tyr Lys Cys Glu 40 45 50			379
aag tgc cac ctg gtg ctg tgc agc ccg aag cag acc gag tgt ggg cac Lys Cys His Leu Val Leu Cys Ser Pro Lys Gln Thr Glu Cys Gly His 55 60 65 70			427
cgc ttc tgc gag agc tgc atg gcg gcc ctg ctg agc tct tca agt cca Arg Phe Cys Glu Ser Cys Met Ala Ala Leu Leu Ser Ser Ser Ser Pro 75 80 85			475
aaa tgt aca gcg tgt caa gag agc atc gtt aaa gat aag gtg ttt aag Lys Cys Thr Ala Cys Gln Glu Ser Ile Val Lys Asp Lys Val Phe Lys 90 95 100			523
gat aat tgc tgc aag aga gaa att ctg gct ctt cag atc tat tgt cgg Asp Asn Cys Cys Lys Arg Glu Ile Leu Ala Leu Gln Ile Tyr Cys Arg 105 110 115			571
aat gaa agc aga ggt tgt gca gag cag tta acg ctg gga cat ctg ctg Asn Glu Ser Arg Gly Cys Ala Glu Gln Leu Thr Leu Gly His Leu Leu 120 125 130			619
gtg cat tta aaa aat gat tgc cat ttt gaa gaa ctt cca tgt gtg cgt Val His Leu Lys Asn Asp Cys His Phe Glu Glu Leu Pro Cys Val Arg 135 140 145 150			667
cct gac tgc aaa gaa aag gtc ttg agg aaa gac ctg cga gac cac gtg Pro Asp Cys Lys Glu Lys Val Leu Arg Lys Asp Leu Arg Asp His Val 155 160 165			715
gag aag gcg tgt aaa tac cgg gaa gcc aca tgc agc cac tgc aag agt Glu Lys Ala Cys Lys Tyr Arg Glu Ala Thr Cys Ser His Cys Lys Ser 170 175 180			763
cag gtt ccg atg atc gcg ctg cag aaa cac gaa gac acc gac tgt ccc Gln Val Pro Met Ile Ala Leu Gln Lys His Glu Asp Thr Asp Cys Pro 185 190 195			811
tgc gtg gtg gtc tcc tgc cct cac aag tgc agc gtc cag act ctc ctg Cys Val Val Val Ser Cys Pro His Lys Cys Ser Val Gln Thr Leu Leu 200 205 210			859
agg agc gag ttg agt gca cac ttg tca gag tgt gtc aat gcc ccc agc Arg Ser Glu Leu Ser Ala His Leu Ser Glu Cys Val Asn Ala Pro Ser 215 220 225 230			907
acc tgt agt ttt aag cgc tat ggc tgc gtt ttt cag ggg aca aac cag Thr Cys Ser Phe Lys Arg Tyr Gly Cys Val Phe Gln Gly Thr Asn Gln 235 240 245			955
cag atc aag gcc cac gag gcc agc tcc gcc gtg cag cac gtc aac ctg Gln Ile Lys Ala His Glu Ala Ser Ser Ala Val Gln His Val Asn Leu 250 255 260			1003
ctg aag gag tgg agc aac tcg ctc gaa aag aag gtt tcc ttg ttg cag Leu Lys Glu Trp Ser Asn Ser Leu Glu Lys Lys Val Ser Leu Leu Gln 265 270 275			1051
aat gaa agt gta gaa aaa aac aag agc ata caa agt ttg cac aat cag			1099

Asn	Glu	Ser	Val	Glu	Lys	Asn	Lys	Ser	Ile	Gln	Ser	Leu	His	Asn	Gln		
280						285				290							
ata	tgt	agc	ttt	gaa	att	gaa	att	gag	aga	caa	aag	gaa	atg	ctt	cga	1147	
Ile	Cys	Ser	Phe	Glu	Ile	Glu	Ile	Glu	Arg	Gln	Lys	Glu	Met	Leu	Arg		
295				300				305						310			
aat	aat	gaa	tcc	aaa	atc	ctt	cat	tta	cag	cga	gtg	atc	gac	agc	caa	1195	
Asn	Asn	Glu	Ser	Lys	Ile	Leu	His	Leu	Gln	Arg	Val	Ile	Asp	Ser	Gln		
				315				320						325			
gca	gag	aaa	ctg	aag	gag	ctt	gac	aag	gag	atc	cgg	ccc	ttc	cgg	cag	1243	
Ala	Glu	Lys	Leu	Lys	Glu	Leu	Asp	Lys	Glu	Ile	Arg	Pro	Phe	Arg	Gln		
			330					335						340			
aac	tgg	gag	gaa	gca	gac	agc	atg	aag	agc	agc	gtg	gag	tcc	ctc	cag	1291	
Asn	Trp	Glu	Glu	Ala	Asp	Ser	Met	Lys	Ser	Ser	Val	Glu	Ser	Leu	Gln		
		345					350						355				
aac	cgc	gtg	acc	gag	ctg	gag	agc	gtg	gac	aag	agt	gcg	ggg	caa	gtg	1339	
Asn	Arg	Val	Thr	Glu	Leu	Glu	Ser	Val	Asp	Lys	Ser	Ala	Gly	Gln	Val		
	360					365					370						
gct	cgg	aac	aca	ggc	ctg	ctg	gag	tcc	cag	ctg	agc	cgg	cat	gac	cag	1387	
Ala	Arg	Asn	Thr	Gly	Leu	Leu	Glu	Ser	Gln	Leu	Ser	Arg	His	Asp	Gln		
375				380						385					390		
atg	ctg	agt	gtg	cac	gac	atc	cgc	cta	gcc	gac	atg	gac	ctg	cgc	ttc	1435	
Met	Leu	Ser	Val	His	Asp	Ile	Arg	Leu	Ala	Asp	Met	Asp	Leu	Arg	Phe		
				395				400						405			
cag	gtc	ctg	gag	acc	gcc	agc	tac	aat	gga	gtg	ctc	atc	tgg	aag	att	1483	
Gln	Val	Leu	Glu	Thr	Ala	Ser	Tyr	Asn	Gly	Val	Leu	Ile	Trp	Lys	Ile		
			410					415						420			
cgc	gac	tac	aag	cgg	cgg	aag	cag	gag	gcc	gtc	atg	ggg	aag	acc	ctg	1531	
Arg	Asp	Tyr	Lys	Arg	Arg	Lys	Gln	Glu	Ala	Val	Met	Gly	Lys	Thr	Leu		
		425					430					435					
tcc	ctt	tac	agc	cag	cct	ttc	tac	act	ggc	tac	ttt	ggc	tat	aag	atg	1579	
Ser	Leu	Tyr	Ser	Gln	Pro	Phe	Tyr	Thr	Gly	Tyr	Phe	Gly	Tyr	Lys	Met		
	440					445						450					
tgt	gcc	agg	gtc	tac	ctg	aac	ggg	gac	ggg	atg	ggg	aag	ggg	acg	cac	1627	
Cys	Ala	Arg	Val	Tyr	Leu	Asn	Gly	Asp	Gly	Met	Gly	Lys	Gly	Thr	His		
455					460				465						470		
ttg	tcg	ctg	ttt	ttt	gtc	atc	atg	cgt	gga	gaa	tat	gat	gcc	ctg	ctt	1675	
Leu	Ser	Leu	Phe	Phe	Val	Ile	Met	Arg	Gly	Glu	Tyr	Asp	Ala	Leu	Leu		
				475					480					485			
cct	tgg	ccg	ttt	aag	cag	aaa	gtg	aca	ctc	atg	ctg	atg	gat	cag	ggg	1723	
Pro	Trp	Pro	Phe	Lys	Gln	Lys	Val	Thr	Leu	Met	Leu	Met	Asp	Gln	Gly		
			490					495						500			
tcc	tct	cga	cgt	cat	ttg	gga	gat	gca	ttc	aag	ccc	gac	ccc	aac	agc	1771	
Ser	Ser	Arg	Arg	His	Leu	Gly	Asp	Ala	Phe	Lys	Pro	Asp	Pro	Asn	Ser		
		505					510					515					
agc	agc	ttc	aag	aag	ccc	act	gga	gag	atg	aat	atc	gcc	tct	ggc	tgc	1819	
Ser	Ser	Phe	Lys	Lys	Pro	Thr	Gly	Glu	Met	Asn	Ile	Ala	Ser	Gly	Cys		
	520					525						530					

cca gtc ttt gtg gcc caa act gtt cta gaa aat ggg aca tat att aaa 1867
 Pro Val Phe Val Ala Gln Thr Val Leu Glu Asn Gly Thr Tyr Ile Lys
 535 540 545 550

gat gat aca att ttt att aaa gtc ata gtg gat act tcg gat ctg ccc 1915
 Asp Asp Thr Ile Phe Ile Lys Val Ile Val Asp Thr Ser Asp Leu Pro
 555 560 565

gat ccc tga taagtagctg gggaggtgga ttagcagaa ggcaactcct 1964
 Asp Pro

ctgggggatt tgaaccgggc tgtcttcact gaggtcctcg cgctcagaaa aggaccttgt 2024

gagacggagg aagcggcaga aggcggacgc gtgccggcgg gaggagccac gcgtgagcac 2084

acctgacacg ttttataata gactagccc acttcactct gaagaattat ttatccttca 2144

acaagataaa tattgctgtc agagaagggt ttcattttca tttttaaaga tctagttaat 2204

taaggtggaa aacatatatg ctaaacaaaa gaaacatgat tttcttcct taaacttgaa 2264

caccaaaaaa acacacacac acacacacgt ggggatagct ggacatgtca gcatgttaag 2324

taaaaggaga atttatgaaa tagtaatgca attctgatat cttctttcta aaattcaaga 2384

gtgcaatttt gtttcaaata cagtatattg tctattttta aggcctccaa aaaaaaaaaa 2444

aattccggcc g 2455

<210> 4
 <211> 1999
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (86)..(1498)

<300>
 <308> X80200
 <309> 1998-04-15

<400> 4

gccgggagcg ccgctccagc gaggcgcggg ctgtggggcc gccgcgtgcc tggccccgct 60

cgcccggtgcc ggccgctcgc ccgcc atg cct ggc ttc gac tac aag ttc ctg 112
 Met Pro Gly Phe Asp Tyr Lys Phe Leu
 1 5

gag aag ccc aag cga cgg ctg ctg tgc cca ctg tgc ggg aag ccc atg 160
 Glu Lys Pro Lys Arg Arg Leu Leu Cys Pro Leu Cys Gly Lys Pro Met
 10 15 20 25

cgc gag cct gtg cag gtt tcc acc tgc ggc cac cgt ttc tgc gat acc 208
 Arg Glu Pro Val Gln Val Ser Thr Cys Gly His Arg Phe Cys Asp Thr
 30 35 40

tgc ctg cag gag ttc ctc agt gaa gga gtc ttc aag tgc cct gag gac 256
 Cys Leu Gln Glu Phe Leu Ser Glu Gly Val Phe Lys Cys Pro Glu Asp
 45 50 55

cag ctt cct ctg gac tat gcc aag atc tac cca gac ccg gag ctg gaa 304

Gln	Leu	Pro	Leu	Asp	Tyr	Ala	Lys	Ile	Tyr	Pro	Asp	Pro	Glu	Leu	Glu		
		60					65					70					
gta	caa	gta	ttg	ggc	ctg	cct	atc	cgc	tgc	atc	cac	agt	gag	gag	ggc	352	
Val	Gln	Val	Leu	Gly	Leu	Pro	Ile	Arg	Cys	Ile	His	Ser	Glu	Glu	Gly		
	75						80				85						
tgc	cgc	ttg	agt	ggg	cca	cta	cgt	cat	cta	cag	ggc	cac	ctg	aat	acc	400	
Cys	Arg	Trp	Ser	Gly	Pro	Leu	Arg	His	Leu	Gln	Gly	His	Leu	Asn	Thr		
	90				95					100					105		
tgc	agc	ttc	aat	gtc	att	ccc	tgc	cct	aat	cgc	tgc	ccc	atg	aag	ctg	448	
Cys	Ser	Phe	Asn	Val	Ile	Pro	Cys	Pro	Asn	Arg	Cys	Pro	Met	Lys	Leu		
				110					115					120			
agc	cgc	cgt	gat	cta	cct	gca	cac	ttg	cag	cat	gac	tgc	ccc	aag	cgg	496	
Ser	Arg	Arg	Asp	Leu	Pro	Ala	His	Leu	Gln	His	Asp	Cys	Pro	Lys	Arg		
			125					130					135				
cgc	ctc	aag	tgc	gag	ttt	tgt	ggc	tgt	gac	ttc	agt	ggg	gag	gcc	tat	544	
Arg	Leu	Lys	Cys	Glu	Phe	Cys	Gly	Cys	Asp	Phe	Ser	Gly	Glu	Ala	Tyr		
		140					145					150					
gag	agc	cat	gag	ggc	atg	tgc	ccc	cag	gag	agt	gtc	tac	tgt	gag	aat	592	
Glu	Ser	His	Glu	Gly	Met	Cys	Pro	Gln	Glu	Ser	Val	Tyr	Cys	Glu	Asn		
	155					160					165						
aag	tgt	ggc	gcc	cgc	atg	atg	cgg	ggg	ctg	ctg	gcc	cag	cat	gcc	acc	640	
Lys	Cys	Gly	Ala	Arg	Met	Met	Arg	Gly	Leu	Leu	Ala	Gln	His	Ala	Thr		
	170				175				180					185			
tct	gag	tgc	ccc	aag	cgc	act	cag	ccc	tgc	acc	tac	tgc	act	aag	gag	688	
Ser	Glu	Cys	Pro	Lys	Arg	Thr	Gln	Pro	Cys	Thr	Tyr	Cys	Thr	Lys	Glu		
			190						195					200			
ttc	gtc	ttt	gac	acc	atc	cag	agc	cac	cag	tac	cag	tgc	cca	agg	ctg	736	
Phe	Val	Phe	Asp	Thr	Ile	Gln	Ser	His	Gln	Tyr	Gln	Cys	Pro	Arg	Leu		
			205					210					215				
cct	gtt	gcc	tgc	ccc	aac	caa	tgt	ggc	gtg	ggc	act	gtg	gct	cgg	gag	784	
Pro	Val	Ala	Cys	Pro	Asn	Gln	Cys	Gly	Val	Gly	Thr	Val	Ala	Arg	Glu		
		220				225						230					
gac	ctg	cca	ggc	cat	ctg	aag	gac	agc	tgt	aac	acc	gcc	ctg	gtg	ctc	832	
Asp	Leu	Pro	Gly	His	Leu	Lys	Asp	Ser	Cys	Asn	Thr	Ala	Leu	Val	Leu		
	235					240					245						
tgc	cca	ttc	aaa	gac	tcc	ggc	tgc	aag	cac	agg	tgc	cct	aag	ctg	gca	880	
Cys	Pro	Phe	Lys	Asp	Ser	Gly	Cys	Lys	His	Arg	Cys	Pro	Lys	Leu	Ala		
	250				255					260				265			
atg	gca	cgg	cat	gtg	gag	gag	agt	gtg	aag	cca	cat	ctg	gcc	atg	atg	928	
Met	Ala	Arg	His	Val	Glu	Glu	Ser	Val	Lys	Pro	His	Leu	Ala	Met	Met		
				270					275					280			
tgt	gcc	ctg	gtg	agc	cgg	caa	cgg	cag	gag	ctg	cag	gag	ctt	cgg	cga	976	
Cys	Ala	Leu	Val	Ser	Arg	Gln	Arg	Gln	Glu	Leu	Gln	Glu	Leu	Arg	Arg		
			285					290					295				
gag	ctg	gag	gag	cta	tca	gtg	ggc	agt	gat	ggc	gtg	ctc	atc	tgg	aag	1024	
Glu	Leu	Glu	Glu	Leu	Ser	Val	Gly	Ser	Asp	Gly	Val	Leu	Ile	Trp	Lys		
		300					305					310					

att ggc agc tat gga cgg cgg cta cag gag gcc aag gcc aag ccc aac 1072
 Ile Gly Ser Tyr Gly Arg Arg Leu Gln Glu Ala Lys Ala Lys Pro Asn
 315 320 325

ctt gag tgc ttc agc cca gcc ttc tac aca cat aag tat ggt tac aag 1120
 Leu Glu Cys Phe Ser Pro Ala Phe Tyr Thr His Lys Tyr Gly Tyr Lys
 330 335 340 345

ctg cag gtg tct gca ttc ctc aat ggc aat ggc agt ggt gag ggc aca 1168
 Leu Gln Val Ser Ala Phe Leu Asn Gly Asn Gly Ser Gly Glu Gly Thr
 350 355 360

cac ctc tca ctg tac att cgt gtg ctg cct ggt gcc ttt gac aat ctc 1216
 His Leu Ser Leu Tyr Ile Arg Val Leu Pro Gly Ala Phe Asp Asn Leu
 365 370 375

ctt gag tgg ccc ttt gcc cgc cgt gtc acc ttc tcc ctg ctg gat cag 1264
 Leu Glu Trp Pro Phe Ala Arg Arg Val Thr Phe Ser Leu Leu Asp Gln
 380 385 390

agc gac cct ggg ctg gct aaa cca cag cac gtc act gag acc ttc cac 1312
 Ser Asp Pro Gly Leu Ala Lys Pro Gln His Val Thr Glu Thr Phe His
 395 400 405

ccc gac cca aac tgg aag aat ttc cag aag cca gcc acg tgg cgg ggc 1360
 Pro Asp Pro Asn Trp Lys Asn Phe Gln Lys Pro Gly Thr Trp Arg Gly
 410 415 420 425

tcc ctg gat gag agt tct ctg ggc ttt ggt tat ccc aag ttc atc tcc 1408
 Ser Leu Asp Glu Ser Ser Leu Gly Phe Gly Tyr Pro Lys Phe Ile Ser
 430 435 440

cac cag gac att cga aag cga aac tat gtg cgg gat gat gca gtc ttc 1456
 His Gln Asp Ile Arg Lys Arg Asn Tyr Val Arg Asp Asp Ala Val Phe
 445 450 455

atc cgt gct gct gtt gaa ctg ccc cgg aag atc ctc agc tga 1498
 Ile Arg Ala Ala Val Glu Leu Pro Arg Lys Ile Leu Ser
 460 465 470

gtgcaggtgg ggttcgaggg gaaaggacga tggggcatga cctcagtcag gcactggctg 1558
 aacttgagaga gggggccgga ccccgctcag ctgcttctgc tgcctaggtt ctgttaccac 1618
 atcctccctc cccagccac caccctcagg tgcctccaat tgggtgcttca gccctggccc 1678
 ctgtggggaa caggtcttgg ggtcatgaag ggctggaaac aagtgacccc agggcctgtc 1738
 tcccttcttg ggtaggcag acatgccttg gtgcgggtca cactctacac ggactgaggt 1798
 gcctgctcag gtgctatgtc ccaagagcca taagggggtg ggaattgggg agggagaaaag 1858
 ggtagttcaa agagtctgtc ttgagatctg attttttccc cctttacctg gctgtgcccc 1918
 ctctggttat ttatttcctt agtgccagga gggcacagca ggggagccct gatttttaat 1978
 aaatccggaa ttgtatttat t 1999

<210> 5
 <211> 3993
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (55)..(1728)

<300>
 <308> AB000509
 <309> 1998-03-25

<400> 5

```

gcagcagccg cgccctgcaga ccggcctgcg ggagcccgcg cgccgagccc caca atg 57
                                     Met
                                     1

gct tat tca gaa gag cat aaa ggt atg ccc tgt ggt ttc atc cgc cag 105
Ala Tyr Ser Glu Glu His Lys Gly Met Pro Cys Gly Phe Ile Arg Gln
                                     5
                                     10
                                     15

aat tcc ggc aac tcc att tcc ttg gac ttt gag ccc agt ata gag tac 153
Asn Ser Gly Asn Ser Ile Ser Leu Asp Phe Glu Pro Ser Ile Glu Tyr
                                     20
                                     25
                                     30

cag ttt gtg gag cgg ctg gaa gag cgc tac aaa tgt gcc ttc tgc cac 201
Gln Phe Val Glu Arg Leu Glu Glu Arg Tyr Lys Cys Ala Phe Cys His
                                     35
                                     40
                                     45

tcg gtg ctt cac aac ccc cac cag aca gga tgt ggg cac cgc ttc tgc 249
Ser Val Leu His Asn Pro His Gln Thr Gly Cys Gly His Arg Phe Cys
                                     50
                                     55
                                     60
                                     65

cag cac tgc atc ctg tcc ctg aga gaa tta aac aca gtg cca atc tgc 297
Gln His Cys Ile Leu Ser Leu Arg Glu Leu Asn Thr Val Pro Ile Cys
                                     70
                                     75
                                     80

cct gta gat aaa gag gtc atc aaa tct cag gag gtt ttt aaa gac aat 345
Pro Val Asp Lys Glu Val Ile Lys Ser Gln Glu Val Phe Lys Asp Asn
                                     85
                                     90
                                     95

tgt tgc aaa aga gaa gtc ctc aac tta tat gta tat tgc agc aat gct 393
Cys Cys Lys Arg Glu Val Leu Asn Leu Tyr Val Tyr Cys Ser Asn Ala
                                     100
                                     105
                                     110

cct gga tgt aat gcc aag gtt att ctg ggc cgg tac cag gat cac ctt 441
Pro Gly Cys Asn Ala Lys Val Ile Leu Gly Arg Tyr Gln Asp His Leu
                                     115
                                     120
                                     125

cag cag tgc tta ttt caa cct gtg cag tgt tct aat gag aag tgc cgg 489
Gln Gln Cys Leu Phe Gln Pro Val Gln Cys Ser Asn Glu Lys Cys Arg
                                     130
                                     135
                                     140
                                     145

gag cca gtc cta cgg aaa gac ctg aaa gag cat ttg agt gca tcc tgt 537
Glu Pro Val Leu Arg Lys Asp Leu Lys Glu His Leu Ser Ala Ser Cys
                                     150
                                     155
                                     160

cag ttt cga aag gaa aaa tgc ctt tat tgc aaa aag gat gtg gta gtc 585
Gln Phe Arg Lys Glu Lys Cys Leu Tyr Cys Lys Lys Asp Val Val Val
                                     165
                                     170
                                     175

atc aat cta cag aat cat gag gaa aac ttg tgt cct gaa tac cca gta 633
Ile Asn Leu Gln Asn His Glu Glu Asn Leu Cys Pro Glu Tyr Pro Val
                                     180
                                     185
                                     190

ttt tgt ccc aac aat tgt gcg aag att att cta aaa act gag gta gat 681

```

Phe	Cys	Pro	Asn	Asn	Cys	Ala	Lys	Ile	Ile	Leu	Lys	Thr	Glu	Val	Asp		
195						200					205						
gaa cac ctg gct gta tgt cct gaa gct gag caa gac tgt cct ttt aag 729																	
Glu	His	Leu	Ala	Val	Cys	Pro	Glu	Ala	Glu	Gln	Asp	Cys	Pro	Phe	Lys		
210					215					220					225		
cac tat ggc tgt gct gta acg gat aaa cgg agg aac ctg cag caa cat 777																	
His	Tyr	Gly	Cys	Ala	Val	Thr	Asp	Lys	Arg	Arg	Asn	Leu	Gln	Gln	His		
				230					235					240			
gag cat tca gcc tta cgg gag cac atg cgt ttg gtt tta gaa aag aat 825																	
Glu	His	Ser	Ala	Leu	Arg	Glu	His	Met	Arg	Leu	Val	Leu	Glu	Lys	Asn		
			245					250					255				
gtc caa tta gaa gaa cag att tct gac tta cac aag agc cta gaa cag 873																	
Val	Gln	Leu	Glu	Glu	Gln	Ile	Ser	Asp	Leu	His	Lys	Ser	Leu	Glu	Gln		
			260			265						270					
aaa gaa agt aaa atc cag cag cta gca gaa act ata aag aaa ctt gaa 921																	
Lys	Glu	Ser	Lys	Ile	Gln	Gln	Leu	Ala	Glu	Thr	Ile	Lys	Lys	Leu	Glu		
	275					280					285						
aag gag ttc aag cag ttt gca cag ttg ttt ggc aaa aat gga agc ttc 969																	
Lys	Glu	Phe	Lys	Gln	Phe	Ala	Gln	Leu	Phe	Gly	Lys	Asn	Gly	Ser	Phe		
290					295					300					305		
ctc cca aac atc cag gtt ttt gcc agt cac att gac aag tca gct tgg 1017																	
Leu	Pro	Asn	Ile	Gln	Val	Phe	Ala	Ser	His	Ile	Asp	Lys	Ser	Ala	Trp		
				310					315					320			
cta gaa gct caa gtg cat caa tta tta caa atg gtt aac cag caa caa 1065																	
Leu	Glu	Ala	Gln	Val	His	Gln	Leu	Leu	Gln	Met	Val	Asn	Gln	Gln	Gln		
			325					330					335				
aat aaa ttt gac ctg aga cct ttg atg gaa gca gtt gat aca gtg aaa 1113																	
Asn	Lys	Phe	Asp	Leu	Arg	Pro	Leu	Met	Glu	Ala	Val	Asp	Thr	Val	Lys		
		340					345					350					
cag aaa att acc ctg cta gaa aac aat gat caa aga tta gcc gtt tta 1161																	
Gln	Lys	Ile	Thr	Leu	Leu	Asn	Asn	Asp	Gln	Arg	Leu	Ala	Val	Leu			
	355					360					365						
gaa gag gaa act aac aaa cat gat acc cac att aat att cat aaa gca 1209																	
Glu	Glu	Glu	Thr	Asn	Lys	His	Asp	Thr	His	Ile	Asn	Ile	His	Lys	Ala		
370						375					380				385		
cag ctg agt aaa aat gaa gag cga ttt aaa ctg ctg gag ggt act tgc 1257																	
Gln	Leu	Ser	Lys	Asn	Glu	Glu	Arg	Phe	Lys	Leu	Leu	Glu	Gly	Thr	Cys		
			390						395					400			
tat aat gga aag ctc att tgg aag gtg aca gat tac aag atg aag aag 1305																	
Tyr	Asn	Gly	Lys	Leu	Ile	Trp	Lys	Val	Thr	Asp	Tyr	Lys	Met	Lys	Lys		
			405					410					415				
aga gag gcg gtg gat ggg cac aca gtg tcc atc ttc agc cag tcc ttc 1353																	
Arg	Glu	Ala	Val	Asp	Gly	His	Thr	Val	Ser	Ile	Phe	Ser	Gln	Ser	Phe		
			420				425					430					
tac acc agc cgc tgt ggc tac cgg ctc tgt gct aga gca tac ctg aat 1401																	
Tyr	Thr	Ser	Arg	Cys	Gly	Tyr	Arg	Leu	Cys	Ala	Arg	Ala	Tyr	Leu	Asn		
	435					440					445						

```

ggg gat ggg tca ggg agg ggg tca cac ctg tcc cta tac ttt gtg gtc 1449
Gly Asp Gly Ser Gly Arg Gly Ser His Leu Ser Leu Tyr Phe Val Val
450 455 460 465

atg cga gga gag ttt gac tca ctg ttg cag tgg cca ttc agg cag agg 1497
Met Arg Gly Glu Phe Asp Ser Leu Leu Gln Trp Pro Phe Arg Gln Arg
470 475 480

gtg acc ctg atg ctt ctg gac cag agt ggc aaa aag aac att atg gag 1545
Val Thr Leu Met Leu Leu Asp Gln Ser Gly Lys Lys Asn Ile Met Glu
485 490 495

acc ttc aaa cct gac ccc aat agc agc agc ttt aaa aga cct gat ggg 1593
Thr Phe Lys Pro Asp Pro Asn Ser Ser Ser Phe Lys Arg Pro Asp Gly
500 505 510

gag atg aac att gca tct ggc tgt ccc cgc ttt gtg gct cat tct gtt 1641
Glu Met Asn Ile Ala Ser Gly Cys Pro Arg Phe Val Ala His Ser Val
515 520 525

ttg gag aat gcc aag aac gcc tac att aaa gat gac act ctg ttc ttg 1689
Leu Glu Asn Ala Lys Asn Ala Tyr Ile Lys Asp Asp Thr Leu Phe Leu
530 535 540 545

aaa gtg gcc gtg gac tta act gac ctg gag gat ctc tag tcaactgttat 1738
Lys Val Ala Val Asp Leu Thr Asp Leu Glu Asp Leu
550 555

gggggtgataa gaggacttct tggggccaga actgtggagg agagcacatt tgattatcat 1798
attgacctgg atttagactc aaagcacatt tgtatttgcc tttttcctta acgtttgaag 1858
tcagtttaaa acttctgaag tgctgtcttt ttacatttta ctctgtccca gtttgaaact 1918
taaaaactctt agaattattct cttattattt atattttttat atttcttgaa agatggtaag 1978
tttcttgaag tttttggggc gtttctcttt tactggtgct tagcgcagtg tctcgggcac 2038
tctaaatatt gagtgttatg gaggacacag aggtagcaga atcccagttg aaaatgtttt 2098
gatattttat tgtttggcct attgattcta gacctggcct taagtctgca aaagccatct 2158
ttataaggta ggctgttcca gttaagaagt gggatgatgta gttacaaaga taatatgctc 2218
agtttggacc tttttttcag ttaaatgcta aatatatgaa aattactata cctctaagta 2278
ttttcatgaa attcaccagc agtttgcaag cacagttttg caaggctgca taagaactgg 2338
tgaatggggg aagcattttc attcttctctg ctgaagtaaa gcagaaagta ctccatagta 2398
tatgagatat agccagctag ctaaagtcca gattttgtta ggttcaacct tatgaaaaaa 2458
actattttca taggtcaaaa atggtaaaaa attagcagtt tcataagatt caaccaaata 2518
aatatatata tacacacaca catacatata cacctatata tgtgtgtata caaacagttc 2578
gaatgtatgt tgggtgacagt aataaatcaa tgtgaggatg gatagaattt agtatatgat 2638
agagaaaatg tcataaatgg ataaaaggaa tttacaactt gaggagaaaa cctttacaat 2698
ttcctatggg tgtcagaagt actctcagcg aaaactgatg gctaaaacag tatctactat 2758
tctctgataa cttttttttt gagacagagt ttcattgtca cccaggctgg agtacagtgg 2818

```

catgatctca gctcactgca aactctgcct cccgaattca agtgattctc ctgcctcagc 2878
ctectgagta gctgggatta caggcgcccc tcaccacacc caggtaattt ttgtattttt 2938
agtagagacg gagttttgcc atgttgGCCa agctgatctc aaactcctga cctcaagtga 2998
tctgcccgcc tgggctctcc aaagtgcctga gattacaggc atgaccaccc gcgtcaagcc 3058
tctgacaact attgaatttg taagctgcta tgcaaattgg catttatata aacttggat 3118
gtttcttgct agaattctga gtactctgtg aagaacagaa atgatcatat tcttatgcat 3178
ctatctgtat gggctcgaag gtgtatatac aaactgagat gagtccttat gactcttgat 3238
aagcctgagt ttaacaacaa caaaaatgcc aagttgtcct gagcccttct gcgttggtat 3298
gccacttccc tactgctcat atgcacgctg gctccctgg gcacgcaagg atgagtatgg 3358
gccatgggcc cctgtagagc tgcttacctg gtgatgacca tgcaccttac aatttctgaa 3418
cagttaaccc tatagaagca tgctttatat gagggtcttc tgggaagagg aaccttctta 3478
atctctcttg tgggattttc aaaatgctaa agactcacac tgcagcaatc atcccagatg 3538
attaaattca aagaaatagg ttcacaacag gaatatactg aagaactaga gtgtcactgc 3598
tgggtgaactg tggcacgggt gctcaacaca tcacctcgga caaattcagg aagcatttct 3658
ttagcccaca agtccagacc cagggtcctt gtatgtttgt ttttaattt catcatatcc 3718
aagttcactc tgtcttctg agcagtgga gatcatattg ctgtaacttc ttttaagtag 3778
ttgatgtgga aaacatttta aagtgaattt gtcaaatgc tgggttttg ttttatccaa 3838
cttttgtgca tatatataaa gtatgtcatg gcatgggttg cttaggagtt cagagttcct 3898
tcatcatcga aatagtgatt aagtgatccc agaacaagga atactagagt aaaaagcacc 3958
tctttttcag aaaaaaaaa aaaaaaaaa aaaaa 3993

<210> 6
<211> 2264
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (222) .. (1790)

<300>
<308> U78798
<309> 1996-12-12

<400> 6

ccgcagctgg ggcttggcct gcgggcggcc agcgaagggtg gcgaaggctc ccaactggatc 60
cagagtttgc cgtccaagca gcctcgtctc ggcgcgaggt gtctgtgtcc gtcctctacc 120
agcgcccttg ctgagcggag tcgtgcgggt ggtgggggag cctgcctc ctggttcggc 180
ctccccgcgc actagaacga gcaagtgata atcaagttac t atg agt ctg cta aac 236
Met Ser Leu Leu Asn

1														5		
tgt	gaa	aac	agc	tgt	gga	tcc	agc	cag	tct	gaa	agt	gac	tgc	tgt	gtg	284
Cys	Glu	Asn	Ser	Cys	Gly	Ser	Ser	Gln	Ser	Glu	Ser	Asp	Cys	Cys	Val	
				10					15					20		
gcc	atg	gcc	agc	tcc	tgt	agc	gct	gta	aca	aaa	gat	gat	agt	gtg	ggg	332
Ala	Met	Ala	Ser	Ser	Cys	Ser	Ala	Val	Thr	Lys	Asp	Asp	Ser	Val	Gly	
			25					30					35			
gga	act	gcc	agc	acg	ggg	aac	ctc	tcc	agc	tca	ttt	atg	gag	gag	atc	380
Gly	Thr	Ala	Ser	Thr	Gly	Asn	Leu	Ser	Ser	Ser	Phe	Met	Glu	Glu	Ile	
			40			45						50				
cag	gga	tat	gat	gta	gag	ttt	gac	cca	ccc	ctg	gaa	agc	aag	tat	gaa	428
Gln	Gly	Tyr	Asp	Val	Glu	Phe	Asp	Pro	Pro	Leu	Glu	Ser	Lys	Tyr	Glu	
			55			60					65					
tgc	ccc	atc	tgc	ttg	atg	gca	tta	cga	gaa	gca	gtg	caa	acg	cca	tgc	476
Cys	Pro	Ile	Cys	Leu	Met	Ala	Leu	Arg	Glu	Ala	Val	Gln	Thr	Pro	Cys	
	70				75				80						85	
ggc	cat	agg	ttc	tgc	aaa	gcc	tgc	atc	ata	aaa	tca	ata	agg	gat	gca	524
Gly	His	Arg	Phe	Cys	Lys	Ala	Cys	Ile	Ile	Lys	Ser	Ile	Arg	Asp	Ala	
				90				95						100		
ggg	cac	aaa	tgt	cca	gtt	gac	aat	gaa	ata	ctg	ctg	gaa	aat	caa	cta	572
Gly	His	Lys	Cys	Pro	Val	Asp	Asn	Glu	Ile	Leu	Leu	Glu	Asn	Gln	Leu	
			105				110						115			
ttt	cca	gac	aat	ttt	gca	aaa	cgt	gag	att	ctt	tct	ctg	atg	gtg	aaa	620
Phe	Pro	Asp	Asn	Phe	Ala	Lys	Arg	Glu	Ile	Leu	Ser	Leu	Met	Val	Lys	
		120					125					130				
tgt	cca	aat	gaa	ggg	tgt	ttg	cac	aag	atg	gaa	ctg	aga	cat	ctt	gag	668
Cys	Pro	Asn	Glu	Gly	Cys	Leu	His	Lys	Met	Glu	Leu	Arg	His	Leu	Glu	
	135					140					145					
gat	cat	caa	gca	cat	tgt	gag	ttt	gct	ctt	atg	gat	tgt	ccc	caa	tgc	716
Asp	His	Gln	Ala	His	Cys	Glu	Phe	Ala	Leu	Met	Asp	Cys	Pro	Gln	Cys	
	150				155					160					165	
cag	cgt	ccc	ttc	caa	aaa	ttc	cat	att	aat	att	cac	att	ctg	aag	gat	764
Gln	Arg	Pro	Phe	Gln	Lys	Phe	His	Ile	Asn	Ile	His	Ile	Leu	Lys	Asp	
			170					175						180		
tgt	cca	agg	aga	cag	gtt	tct	tgt	gac	aac	tgt	gct	gca	tca	atg	gca	812
Cys	Pro	Arg	Arg	Gln	Val	Ser	Cys	Asp	Asn	Cys	Ala	Ala	Ser	Met	Ala	
			185					190					195			
ttt	gaa	gat	aaa	gag	atc	cat	gac	cag	aac	tgt	cct	ttg	gca	aat	gtc	860
Phe	Glu	Asp	Lys	Glu	Ile	His	Asp	Gln	Asn	Cys	Pro	Leu	Ala	Asn	Val	
		200					205					210				
atc	tgt	gaa	tac	tgc	aat	act	ata	ctc	atc	aga	gaa	cag	atg	cct	aat	908
Ile	Cys	Glu	Tyr	Cys	Asn	Thr	Ile	Leu	Ile	Arg	Glu	Gln	Met	Pro	Asn	
	215					220					225					
cat	tat	gat	cta	gac	tgc	cct	aca	gcc	cca	att	cca	tgc	aca	ttc	agt	956
His	Tyr	Asp	Leu	Asp	Cys	Pro	Thr	Ala	Pro	Ile	Pro	Cys	Thr	Phe	Ser	
	230				235					240					245	
act	ttt	ggg	tgc	cat	gaa	aag	atg	cag	agg	aat	cac	ttg	gca	cgc	cac	1004

Thr	Phe	Gly	Cys	His	Glu	Lys	Met	Gln	Arg	Asn	His	Leu	Ala	Arg	His		
				250					255					260			
cta	caa	gag	aac	acc	cag	tca	cac	atg	aga	atg	ttg	gcc	cag	gct	gtt	1052	
Leu	Gln	Glu	Asn	Thr	Gln	Ser	His	Met	Arg	Met	Leu	Ala	Gln	Ala	Val		
			265					270					275				
cat	agt	ttg	agc	gtt	ata	ccc	gac	tct	ggg	tat	atc	tca	gag	gtc	cgg	1100	
His	Ser	Leu	Ser	Val	Ile	Pro	Asp	Ser	Gly	Tyr	Ile	Ser	Glu	Val	Arg		
		280					285					290					
aat	ttc	cag	gaa	act	att	cac	cag	tta	gag	ggg	cgc	ctt	gta	aga	caa	1148	
Asn	Phe	Gln	Glu	Thr	Ile	His	Gln	Leu	Glu	Gly	Arg	Leu	Val	Arg	Gln		
	295					300					305						
gac	cat	caa	atc	cgg	gag	ctg	act	gct	aaa	atg	gaa	act	cag	agt	atg	1196	
Asp	His	Gln	Ile	Arg	Glu	Leu	Thr	Ala	Lys	Met	Glu	Thr	Gln	Ser	Met		
310					315					320					325		
tat	gta	agt	gag	ctc	aaa	cga	acc	att	cga	acc	ctt	gag	gac	aaa	gtt	1244	
Tyr	Val	Ser	Glu	Leu	Lys	Arg	Thr	Ile	Arg	Thr	Leu	Glu	Asp	Lys	Val		
				330					335					340			
gct	gaa	atc	gaa	gca	cag	cag	tgc	aat	gga	att	tat	att	tgg	aag	att	1292	
Ala	Glu	Ile	Glu	Ala	Gln	Gln	Cys	Asn	Gly	Ile	Tyr	Ile	Trp	Lys	Ile		
			345					350					355				
ggc	aac	ttt	gga	atg	cat	ttg	aaa	tgt	caa	gaa	gag	gag	aaa	cct	gtt	1340	
Gly	Asn	Phe	Gly	Met	His	Leu	Lys	Cys	Gln	Glu	Glu	Glu	Lys	Pro	Val		
		360					365					370					
gtg	att	cat	agc	cct	gga	ttc	tac	act	ggc	aaa	ccc	ggg	tac	aaa	ctg	1388	
Val	Ile	His	Ser	Pro	Gly	Phe	Tyr	Thr	Gly	Lys	Pro	Gly	Tyr	Lys	Leu		
	375					380					385						
tgc	atg	cgc	ttg	cac	ctt	cag	tta	ccg	act	gct	cag	cgc	tgt	gca	aac	1436	
Cys	Met	Arg	Leu	His	Gln	Gln	Leu	Pro	Thr	Ala	Gln	Arg	Cys	Ala	Asn		
390					395					400				405			
tat	ata	tcc	ctt	ttt	gtc	cac	aca	atg	caa	gga	gaa	tat	gac	agc	cac	1484	
Tyr	Ile	Ser	Leu	Phe	Val	His	Thr	Met	Gln	Gly	Glu	Tyr	Asp	Ser	His		
				410					415					420			
ctc	cct	tgg	ccc	ttc	cag	ggg	aca	ata	cgc	ctt	aca	att	ctt	gat	cag	1532	
Leu	Pro	Trp	Pro	Phe	Gln	Gly	Thr	Ile	Arg	Leu	Thr	Ile	Leu	Asp	Gln		
			425					430					435				
tct	gaa	gca	cct	gta	agg	caa	aac	cac	gaa	gag	ata	atg	gat	gcc	aaa	1580	
Ser	Glu	Ala	Pro	Val	Arg	Gln	Asn	His	Glu	Glu	Ile	Met	Asp	Ala	Lys		
		440					445					450					
cca	gag	ctg	ctt	gct	ttc	cag	cga	ccc	aca	atc	cca	cgg	aac	cca	aaa	1628	
Pro	Glu	Leu	Leu	Ala	Phe	Gln	Arg	Pro	Thr	Ile	Pro	Arg	Asn	Pro	Lys		
		455				460					465						
ggg	ttt	ggc	tat	gta	act	ttt	atg	cat	ctg	gaa	gcc	cta	aga	caa	aga	1676	
Gly	Phe	Gly	Tyr	Val	Thr	Phe	Met	His	Leu	Glu	Ala	Leu	Arg	Gln	Arg		
470					475					480				485			
act	ttc	att	aag	gat	gac	aca	tta	tta	gtg	cgc	tgt	gag	gtc	tcc	acc	1724	
Thr	Phe	Ile	Lys	Asp	Asp	Thr	Leu	Leu	Val	Arg	Cys	Glu	Val	Ser	Thr		
				490					495					500			

WO 00/20435

PCT/US99/23171

cgc ttt gac atg ggt agc ctt cgg agg gag ggt ttt cag cca cga agt 1772
 Arg Phe Asp Met Gly Ser Leu Arg Arg Glu Gly Phe Gln Pro Arg Ser
 505 510 515

act gat gca ggg gta tag cttgccctca cttgctcaaa aacaactacc 1820
 Thr Asp Ala Gly Val
 520

tggagaaaac agtgcccttc cttgccctgt tctcaataac atgcaaacaa acaagccacg 1880
 ggaaatatgt aatatctact agtgagtgtt gttagagagg tcaattacta tttcttctg 1940
 ttacaaatga tctgaggcag ttttttctg ggaatccaca cgttccatgc tttttcagaa 2000
 atgttaggcc tgaagtgcct gtggcatgtt gcagcagcta ttttgccagt tagtatacct 2060
 ctttgttgta ctttcttggg cttttgtctt ggtgtatttt attgtcagaa agtccagact 2120
 caagagtact aaacttttaa taataatgga ttttccttaa aacttcagtc tttttgtagt 2180
 attatatgta atatattaaa agtgaaaatc actaccgcct tgaaaaaaaa aaaaaaaaaa 2240
 ctcgaggggg gcccgtagcc aatg 2264

<210> 7
 <211> 18
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> antisense sequence

<400> 7
 tttaagttgc tccagggc 18

<210> 8
 <211> 18
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> antisense sequence

<400> 8
 gccgggcgag gactgctg 18

<210> 9
 <211> 18
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> antisense sequence

<400> 9
 gcagacggtg ggagggca 18

<210> 10
 <211> 18
 <212> DNA
 <213> Artificial Sequence

<220>

<223> antisense sequence

<400> 10
ctgggctcct ttgggtcc 18

<210> 11
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 11
cacagcagag agccctgg 18

<210> 12
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 12
attcctcggg ttctcaga 18

<210> 13
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 13
ccattcctcg ggtctca 18

<210> 14
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 14
cctcgccatt cctcgggt 18

<210> 15
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 15
gatcctcgcc attcctcg 18

<210> 16
<211> 18
<212> DNA

<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 16
agacggcttc ctgggctt 18

<210> 17
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 17
ttgaaggagc agccgaca 18

<210> 18
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 18
ggccttccac tgtttcat 18

<210> 19
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 19
ccacttccac ggctgcct 18

<210> 20
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 20
cgcttggtga cattggtg 18

<210> 21
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 21
cgcatcatac tcccctct 18

WO 00/20435

PCT/US99/23171

<210>	22	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	22	
aggcgtcaat	ggcgtgct	18
<210>	23	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	23	
ggaaggcgtc	aatggcgt	18
<210>	24	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	24	
ggaagaagag	tgggcatc	18
<210>	25	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	25	
cgtaggcgtg	cttgggtg	18
<210>	26	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	26	
gccccgccca	ccctaagt	18
<210>	27	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	

<400> 27
ggagccccgc ccacccta 18

<210> 28
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 28
ctcaggagcc ccgcccac 18

<210> 29
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 29
aagggcaggg catcacag 18

<210> 30
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 30
tttgtgcctt gaggtctt 18

<210> 31
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 31
cacccatctt tgtgcctt 18

<210> 32
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 32
ggcctcccag tgtcgcat 18

<210> 33
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 33
cccggtcctg tttctgac 18

<210> 34
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 34
gcaccccatc ccttccac 18

<210> 35
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 35
tggagccgctc tgggtttg 18

<210> 36
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 36
gtcttcaaatt ccaacccc 18

<210> 37
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 37
ttctgggctg gaaggaaa 18

<210> 38
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 38
actttctggg ctggaagg 18

<210> 39
<211> 18

WO 00/20435

PCT/US99/23171

<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 39
agagactttc tgggctgg 18

<210> 40
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 40
tttccagaac ccctgtag 18

<210> 41
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 41
atgtttccag aacccctg 18

<210> 42
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 42
gggctgggtg tgctcctg 18

<210> 43
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 43
tttatgcccc tcttcttc 18

<210> 44
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 44
ggaaagtta tgccctc 18

<210> 45
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 45
tacgggattc tggaaagc 18

<210> 46
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 46
aggtgttacg ggattctg 18

<210> 47
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 47
gtcgcagcgc gccggaattc 20

<210> 48
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 48
ccaacggtcg cagcgcgccg 20

<210> 49
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 49
cagccatgag agctgtgacc 20

<210> 50
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400>	50	
acgctagctg	cagccatgag	20
<210>	51	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	51	
gccacactgc	gcctggaagg	20
<210>	52	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	52	
ccggcaggct	ctccacctcc	20
<210>	53	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	53	
gcagcggcct	tcgtggcagc	20
<210>	54	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	54	
cctcgtggtg	cgcttcacg	20
<210>	55	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	55	
ctcgacactt	gccacaagtc	20
<210>	56	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	

<220>
<223> antisense sequence

<400> 56
cactgcacct cgtgctcctg 20

<210> 57
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 57
cctctgcagg agctctgacc 20

<210> 58
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 58
cagccggtgc tgccggctgc 20

<210> 59
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 59
ccggtgccgt cggcgttcag 20

<210> 60
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 60
acgtcgggcc tgaaggcgtc 20

<210> 61
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 61
ctgtcaggtc cacaatggcc 20

<210> 62

WO 00/20435

PCT/US99/23171

<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	62	
gccggctgtg cctggctgcc		20
<210>	63	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	63	
cttggctgca ggccgacacc		20
<210>	64	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	64	
cggccaatgc caccacagcc		20
<210>	65	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	65	
actgtgctcc tgctacatgg		20
<210>	66	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	66	
gctctggcca gcaggaggcc		20
<210>	67	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	67	
ccacagccag cctggccaag		20

<210> 68
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 68
ctctgtcttc gtgagctgga 20

<210> 69
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> control sequence

<400> 69
cctcgtgctg cggcttcacg 20

<210> 70
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> control sequence

<400> 70
cctggtgctc cggcttcacg 20

<210> 71
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 71
agagccgacg accgccgc 18

<210> 72
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 72
ggaagagccg acgaccgc 18

<210> 73
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

WO 00/20435

PCT/US99/23171

<400>	73	
cgcgccagga	gagtcctat	18
<210>	74	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	74	
ttagcggcgg	gttagtct	18
<210>	75	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	75	
agctttagcg	gcgggtta	18
<210>	76	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	76	
ctcggctctgc	ttcgggct	18
<210>	77	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	77	
tgcccacact	cggtctgc	18
<210>	78	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	78	
cggtgcccac	actcggtc	18
<210>	79	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	

<220>
<223> antisense sequence

<400> 79
gaagcgggtgc ccacactc 18

<210> 80
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 80
ttacacgcct tcgccacg 18

<210> 81
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 81
gtattttacac gccttctc 18

<210> 82
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 82
ccggtattta cagcctt 18

<210> 83
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 83
gagggcagga caccacca 18

<210> 84
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 84
tgtgagggca ggacacca 18

<210> 85
<211> 18

<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 85
cacttgtag gag ggcaggac 18

<210> 86
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 86
gctgggttgt cccctgaa 18

<210> 87
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 87
atctgctggt ttgtcccc 18

<210> 88
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 88
cgcggttctg gagggact 18

<210> 89
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 89
ccccgcactc ttgtccac 18

<210> 90
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 90
ttgccccgca ctcttgtc 18

<210> 91
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 91
ccacttgccc cgcactct 18

<210> 92
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 92
gagccacttg ccccgcac 18

<210> 93
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 93
ttccgagcca cttgcccc 18

<210> 94
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 94
tccgccgctt gtagtcgc 18

<210> 95
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 95
tgcttccgcc gcttgtag 18

<210> 96
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 96

tccctgcttcc gccgcttg 18

<210> 97
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 97
gtccccgttc aggttagac 18

<210> 98
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 98
tcccgtcccc gttcaggt 18

<210> 99
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 99
ccatcccgtc cccgttca 18

<210> 100
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 100
tccccatccc gtccccgt 18

<210> 101
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 101
cccttcccca tcccgtcc 18

<210> 102
<211> 18
<212> DNA
<213> Artificial Sequence

<220>

<223> antisense sequence

<400> 102
tgcgtcccct tccccatc 18

<210> 103
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 103
aagtgcgtcc ccttcccc 18

<210> 104
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 104
cgacaagtgc gtcccctt 18

<210> 105
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 105
aaggaagcag ggcattcat 18

<210> 106
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 106
ctctccagtg ggcttctt 18

<210> 107
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 107
tcattctctcc agtgggct 18

<210> 108
<211> 18
<212> DNA

<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 108
gctaaatcca cctcccca 18

<210> 109
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 109
tctgccgctt cctccgtc 18

<210> 110
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 110
ccgccttctg ccgcttcc 18

<210> 111
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 111
gcatggcggg cgagcggc 18

<210> 112
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 112
ccgtcgcttg ggcttctc 18

<210> 113
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 113
gggcacttga agactcct 18

<210> 114
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 114
ctcagggcac ttgaagac 18

<210> 115
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 115
tggtcctcag ggcacttg 18

<210> 116
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 116
aagctgggcc tcagggca 18

<210> 117
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 117
gcggcagccc tcctcact 18

<210> 118
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 118
ctccagcggc agccctcc 18

<210> 119
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400>	119	
tagggcaggg	aatgacat	18
<210>	120	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	120	
cgattagggc	agggaatg	18
<210>	121	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	121	
gggcagcgat	tagggcag	18
<210>	122	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	122	
gcctccccac	tgaagtca	18
<210>	123	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	123	
atgcgggcac	cacactta	18
<210>	124	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	124	
gggcaggcaa	caggcagc	18
<210>	125	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	

<220>
<223> antisense sequence

<400> 125
ccacagtgcc cacaccac 18

<210> 126
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 126
cgagccacag tgcccaca 18

<210> 127
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 127
tcctcccgag ccacagtg 18

<210> 128
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 128
caggtcctcc cgagccac 18

<210> 129
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 129
ggcagagcac cagggcgg 18

<210> 130
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 130
ctttgaatgg gcagagca 18

<210> 131
<211> 18

<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 131
ggagtctttg aatgggca 18

<210> 132
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 132
atgccgtgcc attgccag 18

<210> 133
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 133
ctcaccaggg cacacatc 18

<210> 134
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 134
cagctcctgc cgttgccg 18

<210> 135
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 135
atgagcacgc catcactg 18

<210> 136
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 136
tgtagccgcc qtccatag 18

<210> 137
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 137
gcctcctgta gccgccgt 18

<210> 138
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 138
tagaaggctg ggctgaag 18

<210> 139
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 139
gtgtgtagaa ggctgggc 18

<210> 140
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 140
gtgtgccctc accactgc 18

<210> 141
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 141
gacacggcgg gcaaaggg 18

<210> 142
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400>	142	
gaaggtgaca	cggcgggc	18
<210>	143	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	143	
gcccagggtc	gctctgat	18
<210>	144	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	144	
cttccagttt	gggtcggg	18
<210>	145	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	145	
gataaccaaa	gcccagag	18
<210>	146	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	146	
catcgtcctt	tcccctcg	18
<210>	147	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	147	
ggccagggtc	gaagcacc	18
<210>	148	
<211>	18	
<212>	DNA	
<213>	Artificial Sequence	

<220>
<223> antisense sequence

<400> 148
ttgtttccag cccttcat 18

<210> 149
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 149
catgtctgcc ctacccaa 18

<210> 150
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 150
gctccccctgc tgtgccct 18

<210> 151
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 151
tgaataagcc attgtggg 18

<210> 152
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 152
ctttatgctc ttctgaat 18

<210> 153
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 153
ggatgaaacc acagggca 18

<210> 154
<211> 18

<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 154
tcaaagtcca aggaaatg 18

<210> 155
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 155
tgaagcaccg agtggcag 18

<210> 156
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 156
gggcagattg gcactgtg 18

<210> 157
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 157
ctcctgagat ttgatgac 18

<210> 158
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 158
ctttccgtag gactggct 18

<210> 159
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 159

gattctgtag attgatga 18

<210> 160
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 160
ttcatctacc tcagtttt 18

<210> 161
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 161
tccgttacag cacagcca 18

<210> 162
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 162
gcatgtgctc ccgtaagg 18

<210> 163
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 163
cttttcaagt ttctttat 18

<210> 164
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 164
cttccatcaa aggtctca 18

<210> 165
<211> 18
<212> DNA
<213> Artificial Sequence

<220>

<223> antisense sequence

<400> 165
tctaaaacgg ctaatctt 18

<210> 166
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 166
tcattcttgta atctgtca 18

<210> 167
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 167
ggactggctg aagatgga 18

<210> 168
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 168
ccctccctga cccatccc 18

<210> 169
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 169
gaatgagcca caaagcgg 18

<210> 170
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 170
caagaacaga gtgtcatc 18

<210> 171
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 171
gtctaaatcc aggtcaat 18

<210> 172
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 172
aaacttacca tctttcaa 18

<210> 173
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 173
ctctgtgtcc tccataac 18

<210> 174
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 174
cttaactgga acagccta 18

<210> 175
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 175
gcaggaagaa tgaaaatg 18

<210> 176
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 176
tatttggttg aatcttat 18

<210> 177
<211> 18

<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 177
aaattctatc catcctca 18

<210> 178
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 178
aaattgtaaa gggtttct 18

<210> 179
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 179
acaatgaaac tctgtctc 18

<210> 180
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 180
gcaaaaactcc gtctctac 18

<210> 181
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 181
caatagttgt cagaggct 18

<210> 182
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 182
aaggactcat ctcagttt 18

<210> 183
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 183
taacaacgca gaagggt 18

<210> 184
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 184
agtagggaag tggcataa 18

<210> 185
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 185
catcaccagg taagcagc 18

<210> 186
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 186
tcctgttggtg aacctatt 18

<210> 187
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 187
ggacttggtg gctaaaga 18

<210> 188
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 188
gctcaggaag acagagtg 18

<210> 189
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 189
tgaactccta agcaaacc 18

<210> 190
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 190
gatgatgaag gaactctg 18

<210> 191
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 191
aggccaagcc ccagctgcgg 20

<210> 192
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 192
cgccaccctc gctggccgcc 20

<210> 193
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 193
gagacgaggc tgcttggacg 20

<210> 194
<211> 20
<212> DNA
<213> Artificial Sequence
<220>

<223> antisense sequence

<400> 194
ggacacagac actgcgcgcc 20

<210> 195
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 195
ccaaggcgct ggtagaggac 20

<210> 196
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 196
ttgctcggtc tagtgcgcgg 20

<210> 197
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 197
catagtaact tgattatcac 20

<210> 198
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 198
agcagactca tagtaacttg 20

<210> 199
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 199
acagtttagc agactcatag 20

<210> 200
<211> 20
<212> DNA

<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 200
acagcgctac aggagctggc 20

<210> 201
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 201
attgatttta tgatgcaggc 20

<210> 202
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 202
gtgacctgca tcccttattg 20

<210> 203
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 203
gtctcagttc catcttgtgc 20

<210> 204
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 204
agagcaaact cacaatgtgc 20

<210> 205
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 205
ttttggaagg gacgctggca 20

<210> 206
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 206
aaatgccatt gatgcagcac 20

<210> 207
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 207
attcacagat gacatttgcc 20

<210> 208
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 208
cgtgccaagt gattcctctg 20

<210> 209
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 209
ggtgttctct tgtaggtggc 20

<210> 210
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 210
ggccaacatt ctcatgtgtg 20

<210> 211
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 211
cgctcaaact atgaacagcc 20

<210> 212
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 212
aggcgcaccct ctaactggtg 20

<210> 213
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 213
ccatttttagc agtcagctcc 20

<210> 214
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 214
cgaatgggttc gtttgagctc 20

<210> 215
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 215
ccattgcact gctgtgcttc 20

<210> 216
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 216
gcagtcggtg actgaaggtg 20

<210> 217
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 217
gccttacagg tgcttcagac 20

<210> 218
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 218
agcaagcagc tctggtttgg 20

<210> 219
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 219
ggctacccat gtcaaagcgg 20

<210> 220
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 220
ttgtttttga gcaagtggg 20

<210> 221
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 221
ggcactgttt tctccaggta 20

<210> 222
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 222
acatatttcc cgtggcttgt 20

<210> 223
<211> 20

<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 223
ggaacgtgtg gattcccagg 20

<210> 224
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 224
tgctgcaaca tgccacaggc 20

<210> 225
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 225
atacaccaga gcaaaagccc 20

<210> 226
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> control sequence

<400> 226
aaaagactga acttttaagg 20

<210> 227
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> control sequence

<400> 227
acttaattac catgactagt 20

<210> 228
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> control sequence

<400> 228
ccacgaggag caccatcaag 20

<210>	229	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	229	
ggaccagcct	tgtggagtcc	20
<210>	230	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	230	
tcaggggtcc	aggctggcca	20
<210>	231	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	231	
tctcagggtt	ccaggctggc	20
<210>	232	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	232	
catctcaggg	ttccaggctg	20
<210>	233	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	233	
gccatctcag	ggttccaggc	20
<210>	234	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	

<400> 234
aggccatctc aggtttccag 20

<210> 235
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 235
ggaggccatc tcagggttcc 20

<210> 236
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 236
ctggaggcca tctcagggtt 20

<210> 237
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 237
agctggaggc catctcaggg 20

<210> 238
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 238
tgagctggag gccatctcag 20

<210> 239
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 239
cctgagctgg aggccatctc 20

<210> 240
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 240
tgcctgagct ggaggccatc 20

<210> 241
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 241
gctgcctgag ctggaggcca 20

<210> 242
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 242
gggcgaggac tgctgcctga 20

<210> 243
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 243
tctcagagag acagcctgca 20

<210> 244
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 244
tcctgggctt atagactgga 20

<210> 245
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 245
gggcttcctt tgaaggagca 20

<210> 246
<211> 20

<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 246
caacagcagg tttaggtggg 20

<210> 247
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 247
cagggccatg ggcccagact 20

<210> 248
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 248
cccggtagca atcgacctcc 20

<210> 249
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 249
ctcagccaga agcttctcct 20

<210> 250
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 250
ggccagggcc aggtgggagg 20

<210> 251
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 251
tgctgaagct ccaccaccct 20

<210> 252
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 252
gtgatcttcc acaggaaagt 20

<210> 253
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 253
ttggcagtgtagaaggctgg 20

<210> 254
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 254
aagagcgaca gatgggttct 20

<210> 255
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 255
gcagcatgaa ggtgaccttg 20

<210> 256
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 256
tctggaagga cgctgagctt 20

<210> 257
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400>	257	
ctccacaatg	cacttgagga	20
<210>	258	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	258	
tgccctgagg	tcttgggtgc	20
<210>	259	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	259	
atcctaacca	gatggccagc	20
<210>	260	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	260	
actggcctcc	cagtgtcgca	20
<210>	261	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	261	
gcttgggtcc	tacggttcca	20
<210>	262	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	antisense sequence	
<400>	262	
tcctgtttct	gaccctggag	20
<210>	263	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	

<220>
<223> antisense sequence

<400> 263
ctgggtttgc ttgttcacct 20

<210> 264
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 264
ccaggaggct agaattgagag 20

<210> 265
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 265
tgaggagctg ggaggacagg 20

<210> 266
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 266
ttgggaagct gagctgccag 20

<210> 267
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 267
acagtgactg ggtttcacct 20

<210> 268
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense sequence

<400> 268
ttacgggatt ctggaaagca 20

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/23171

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) : C07H 21/04; C12Q 1/68; A61K 48/00 US CL : 435/6; 514/44; 536/23.1, 24.5 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/6; 514/44; 536/23.1, 24.5 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched NONE Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) WEST, SCISEARCH, CAPLUS, BIOSIS, MEDLINE, EMBASE		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	XU et al. A Role for TRAF 2 and TRAF 6 in Cytokine-Mediated Induction of E-selectin. Immunity. 1996. vol. 5, pages 407-415, see entire document.	1-7, 9-14, and 17-19 ----- 8
Y	SANGHVI et al. HETEROCYCLIC BASE MODIFICATIONS IN NUCLEIC ACIDS AND THEIR APPLICATIONS IN ANTISENSE OLIGONUCLEOTIDES. Antisense Research and Applications, pages 273-288, see entire document.	8
A	BRANCH, A good antisense molecule is hard to find. TIBS, February 1998. vol. 23, pages 45-50, see entire document.	1-19
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "B" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "A" document member of the same patent family		
Date of the actual completion of the international search 30 NOVEMBER 1999		Date of mailing of the international search report 04 FEB 2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer ANDREW WANG Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)*